

09/25/98

Use type a plus (+) inside this box → ☐

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PTO/SB/29 (12/97)

Approved for use through 09/30/00. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

# UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.	7682-45	Total Pages	82
First Named Inventor or Application Identifier			
Jin et al.			
Express Mail Label No.	EL 134 447 824 US		

11 U.S. to  
09/16/98  
19/25/98

## APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

- ☒ Fee Transmittal Form  
*Submit an original, and a duplicate for fee processing*
- ☒ Specification (Total Pages 68)  
*(preferred arrangement set forth below)*
  - Descriptive title of the Invention
  - Cross Reference to Related Applications
  - Statement Regarding Fed sponsored R&D
  - Reference to Microfiche Appendix
  - Background of the Invention
  - Brief Summary of the Invention
  - Brief Description of the Drawings *(if filed)*
  - Detailed Description of the Invention (including drawings, if filed)
  - Claim(s)
  - Abstract of the Disclosure
- ☒ Drawing(s) (35 USC 113) (Total Sheets 12)
- ☒ Oath or Declaration (Unexecuted) (Total Sheets 2)
  - ☐ Newly executed (original or copy)
  - ☐ Copy from a prior application (37 CFR 1.63(d))  
*(for continuation/divisional with Box 17 completed)*  
*(Note Box 5 below)*
  - ☐ DELETION OF INVENTOR(S)

Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33 (b).

- ☐ Microfiche Computer Program (Appendix)
- ☐ Nucleotide and/or Amino Acid Sequence Submission  
*(if applicable, all necessary)*
  - ☐ Computer Readable Copy
  - ☐ Paper Copy (identical to computer copy)
  - ☐ Statement verifying identity of above copies

## ACCOMPANYING APPLICATION PARTS

- ☐ Assignment Papers (cover sheet & document(s))
- ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney  
*(when there is an assignee)*
- ☐ English Translation Document *(if applicable)*
- ☐ Information Disclosure ☐ Copies of IDS  
Statement (IDS)/PTO-1449 Citations
- ☐ Preliminary Amendment
- ☐ Return Receipt Postcard (MPEP 503)  
*(Should be specifically itemized)*
- ☐ Small Entity ☐ Statement filed in prior application,  
Statement(s) Status still proper and desired
- ☐ Certified Copy of Priority Document(s)  
*(if foreign priority is claimed)*
- ☐ Other:

- If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:  
☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No:

## 18. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

20583

(Insert Customer No. or Attach bar code label here)

☐ Correspondence address below

NAME			
ADDRESS			
CITY	STATE	ZIP CODE	
COUNTRY	TELEPHONE	FAX	

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

**PENNIE & EDMONDS LLP**

COUNSELLORS AT LAW  
1155 Avenue of the Americas  
New York, N.Y. 10036-2711  
(212) 790-9090

ATTORNEY DOCKET NO. 7682-045Date September 25, 1998

Assistant Commissioner for Patents  
Box PATENT APPLICATION  
Washington, D.C. 20231

Sir:

The following utility patent application is enclosed for filing:

Applicant(s): Hong Jin et al.Executed on: UNEXECUTEDTitle of Invention: RECOMBINANT RSV EXPRESSION SYSTEMS AND VACCINES

Pages of Specification 67 + ABSTRACT

Sheets of Drawings 12**PATENT APPLICATION FEE VALUE**

TYPE	NO. FILED	LESS	EXTRA	EXTRA RATE	FEE
Total Claims	27	-20	7	\$22.00 each	254.00
Independent	7	-3	4	\$80.00 each	320.00
Minimum Fee					790.00
Multiple Dependency Fee If Applicable (\$260.00)					
Total					1,364.00
50 % Reduction for Independent Inventor, Nonprofit Organization or Small Business Concern (a verified statement as to the applicant's status is attached)					-
Total Filing Fee					\$ 1,364.00

Priority of application no. 60/060,153, filed September 26, 1997, application no. 60/084,133 filed May 1, 1998, and application no. 60/089,207, filed June 12, 1998 is claimed under 35 U.S.C. § 119(e).

Please charge the required fee to Pennie & Edmonds LLP Deposit Account No. 16-1150. A copy of this sheet is enclosed.

Respectfully submitted,

Laura A. Coruzzi 30,742  
LAURA A. CORUZZI (Reg. No.)  
PENNIE & EDMONDS LLP

Enclosure

This form is not for use with re-issue, design or plant patent applications.

by Margaret B. Poiraudon  
Reg. No. 40,922

## RECOMBINANT RSV VIRUS EXPRESSION SYSTEMS AND VACCINES

### 1. INTRODUCTION

The present invention relates to recombinant  
5 negative strand virus RNA templates which may be used to  
express heterologous gene products in appropriate host cell  
systems and/or to construct recombinant viruses that express,  
package, and/or present the heterologous gene product. The  
expression products and chimeric viruses may advantageously be  
10 used in vaccine formulations. In particular, the present  
invention relates to methods of generating recombinant  
respiratory syncytial viruses and the use of these recombinant  
viruses as expression vectors and vaccines. The invention is  
described by way of examples in which recombinant respiratory  
15 syncytial viral genomes are used to generate infectious viral  
particles.

### 2. BACKGROUND OF THE INVENTION

A number of DNA viruses have been genetically  
20 engineered to direct the expression of heterologous proteins  
in host cell systems (e.g., vaccinia virus, baculovirus,  
etc.). Recently, similar advances have been made with  
positive-strand RNA viruses (e.g., poliovirus). The  
expression products of these constructs, i.e., the  
25 heterologous gene product or the chimeric virus which  
expresses the heterologous gene product, are thought to be  
potentially useful in vaccine formulations (either subunit or  
whole virus vaccines). One drawback to the use of viruses  
such as vaccinia for constructing recombinant or chimeric  
30 viruses for use in vaccines is the lack of  
variation in its major epitopes. This lack of variability in  
the viral strains places strict limitations on the repeated  
use of chimeric vaccinia, in that multiple vaccinations will  
generate host-resistance to the strain so that the inoculated  
35 virus cannot infect the host. Inoculation of a resistant  
individual with chimeric vaccinia will, therefore, not induce  
immune stimulation.

By contrast, negative-strand RNA viruses such as influenza virus and respiratory syncytial virus, demonstrate a wide variability of their major epitopes. Indeed, thousands of variants of influenza have been identified; each strain  
5 evolving by antigenic drift. The negative-strand viruses such as influenza and respiratory syncytial virus would be attractive candidates for constructing chimeric viruses for use in vaccines because its genetic variability allows for the construction of a vast repertoire of vaccine formulations  
10 which will stimulate immunity without risk of developing a tolerance.

### 2.1. RESPIRATORY SYNCYTIAL VIRUS

Virus families containing enveloped single-stranded  
15 RNA of the negative-sense genome are classified into groups having non-segmented genomes (Paramyxoviridae, Rhabdoviridae) or those having segmented genomes (Orthomyxoviridae, Bunyaviridae and Arenaviridae). Paramyxoviridae have been classified into three genera: paramyxovirus (sendai virus,  
20 parainfluenza viruses types 1-4, mumps, newcastle disease virus); morbillivirus (measles virus, canine distemper virus and rinderpest virus); and pneumovirus (respiratory syncytial virus and bovine respiratory syncytial virus).

Human respiratory syncytial virus (RSV) is the leading  
25 cause of severe lower respiratory tract disease in infants and young children and is responsible for considerable morbidity and mortality. Two antigenically diverse RSV subgroups A and B are present in human populations. RSV is also recognized as an important agent of disease in immuno-compromised adults and  
30 in the elderly. Due to the incomplete resistance to RSV reinfection induced by natural infection, RSV may infect multiple times during childhood and life. The goal of RSV immunoprophylaxis is to induce sufficient resistance to prevent the serious disease which may be associated with RSV  
35 infection. The current strategies for developing RSV vaccines principally revolve around the administration of purified viral antigen or the development of live attenuated RSV for

intranasal administration. However, to date there have been no approved vaccines or highly effective antiviral therapy for RSV.

- Infection with RSV can range from an unnoticeable
- 5 infection to severe pneumonia and death. RSV possesses a single-stranded nonsegmented negative-sense RNA genome of 15,221 nucleotides (Collins, 1991, *In The paramyxoviruses* pp. 103-162, D.W. Kingsbury (ed.) Plenum Press, New York). The genome of RSV encodes 10 mRNAs (Collins et al., 1984, J. Virol. 49: 572-578). The genome contains a 44 nucleotide leader sequence at the 3' termini followed by the NS1-NS2-N-P-M-SH-G-F-M2-L and a 155 nucleotide trailer sequence at the 5' termini (Collins, 1991, *supra*). Each gene transcription unit contains a short stretch of conserved gene start (GS) sequence
- 10 and a gene end (GE) sequences.

- The viral genomic RNA is not infectious as naked RNA. The RNA genome of RSV is tightly encapsidated with the major nucleocapsid (N) protein and is associated with the phosphoprotein (P) and the large (L) polymerase subunit.
- 20 These proteins form the nucleoprotein core, which is recognized as the minimum unit of infectivity (Brown et al., 1967, J. Virol. 1: 368-373). The RSV N, P, and L proteins form the viral RNA dependent RNA transcriptase for transcription and replication of the RSV genome (Yu et al., 1995, J. Virol. 69:2412-2419; Grosfeld et al., 1995, J. Virol. 69:5677-86). Recent studies indicate that the M2 gene products (M2-1 and M2-2) are involved and are required for transcription (Collins et al., 1996, Proc. Natl. Acad. Sci. 93:81-5).
- 30 The M protein is expressed as a peripheral membrane protein, whereas the F and G proteins are expressed as integral membrane proteins and are involved in virus attachment and viral entry into cells. The G and F proteins are the major antigens that elicit neutralizing antibodies in
- 35 vivo (as reviewed in McIntosh and Chanock, 1990 "Respiratory Syncytial Virus" 2nd ed. Virology (D. M. Knipe et al., Ed.) Raven Press, Ltd., N.Y.). Antigenic dimorphism between the

subgroups of RSV A and B is mainly linked to the G glycoprotein, whereas the F glycoprotein is more closely related between the subgroups.

- Despite decades of research, no safe and effective RSV vaccine has been developed for the prevention of severe morbidity and mortality associated with RSV infection. A formalin-inactivated virus vaccine has failed to provide protection against RSV infection and its exacerbated symptoms during subsequent infection by the wild-type virus in infants
- 10 (Kapikian et al., 1969, Am. J. Epidemiol. 89:405-21; Chin et al., 1969, Am. J. Epidemiol. 89:449-63) Efforts since have focused on developing live attenuated temperature-sensitive mutants by chemical mutagenesis or cold passage of the wild-type RSV (Gharpure et al., 1969, J. Virol. 3: 414-21; Crowe et
- 15 al., 1994, Vaccine 12: 691-9). However, earlier trials yielded discouraging results with these live attenuated temperature sensitive mutants. Virus candidates were either underattenuated or overattenuated (Kim et al., 1973, Pediatrics 52:56-63; Wright et al., 1976, J. Pediatrics
- 20 88:931-6) and some of the vaccine candidates were genetically unstable which resulted in the loss of the attenuated phenotype (Hodes et al., 1974, Proc. Soc. Exp. Biol. Med. 145:1158-64).

- Attempts have also been made to engineer recombinant
- 25 vaccinia vectors which express RSV F or G envelope glycoproteins. However, the use of these vectors as vaccines to protect against RSV infection in animal studies has shown inconsistent results (Olmsted et al. 1986, Proc. Natl. Acad. Sci. 83:7462-7466; Collins et al., 1990, Vaccine 8:164-168).

- 30 Thus, efforts have turned to engineering recombinant RSV to generate vaccines. For a long time, negative-sense RNA viruses were refractory to study. Only recently has it been possible to recover negative strand RNA viruses using a recombinant reverse genetics approach (U.S. Patent No.
- 35 5,166,057 to Palese et al.). Although this method was originally applied to engineer influenza viral genomes (Luytjes et al. 1989, Cell 59:1107-1113; Enami et al. 1990,

Proc. Natl. Acad. Sci. USA 92: 11563-11567), it has been successfully applied to a wide variety of segmented and nonsegmented negative strand RNA viruses, including rabies (Schnell et al. 1994, EMBO J. 13: 4195-4203); VSV (Lawson et al., 1995, Proc. Natl. Acad. Sci. USA 92: 4477-81); measles virus (Radecke et al., 1995, EMBO J. 14:5773-84); rinderpest virus (Baron & Barrett, 1997, J. virol. 71: 1265-71); human parainfluenza virus (Hoffman & Banerjee, 1997, J. Virol. 71:3272-7; Dubin et al., 1997, Virology 235:323-32); SV5 (He et al., 1997, Virology 237:249-60); respiratory syncytial virus (Collins et al. 1991, Proc. Natl. Acad. Sci. USA 88: 9663-9667) and Sendai virus (Park et al. 1991, Proc. Natl. Acad. Sci. USA 88:5537-5541; Kato et al. 1996, Genes to Cells 1:569-579). Although this approach has been used to successfully rescue RSV, a number of groups have reported that RSV is still refractory to study given several properties of RSV which distinguish it from the better characterized paramyxoviruses of the genera Paramyxovirus, Rubulavirus, and Morbillivirus. These differences include a greater number of RNAs, an unusual gene order at the 3' end of the genome, extensive strain-to-strain sequence diversity, several proteins not found in other nonsegmented negative strand RNA viruses and a requirement for the M2 protein (ORF1) to proceed with full processing of full length transcripts and rescue of a full length genome (Collins et al. PCT WO97/12032; Collins, P.L. et al. pp 1313-1357 of volume 1, Fields Virology, et al., Eds. (3rd ed., Raven Press, 1996).

### 3. SUMMARY OF THE INVENTION

The present invention relates to genetically engineered recombinant RS viruses and viral vectors which contain heterologous genes which for the use as vaccines. In accordance with the present invention, the recombinant RS viral vectors and viruses are engineered to contain heterologous genes, including genes of other viruses, pathogens, cellular genes, tumor antigens, or to encode combinations of genes from different strains of RSV.

Recombinant negative-strand viral RNA templates are described which may be used to transfect transformed cell that express the RNA dependent RNA polymerase and allow for complementation. Alternatively, a plasmid expressing the components of the RNA polymerase from an appropriate promoter can be used to transfect cells to allow for complementation of the negative-strand viral RNA templates. Complementation may also be achieved with the use of a helper virus or wild-type virus to provide the RNA dependent RNA polymerase. The RNA templates are prepared by transcription of appropriate DNA sequences with a DNA-directed RNA polymerase. The resulting RNA templates are of negative-or positive-polarity and contain appropriate terminal sequences which enable the viral RNA-synthesizing apparatus to recognize the template. Bicistronic mRNAs can be constructed to permit internal initiation of translation of viral sequences and allow for the expression of foreign protein coding sequences from the regular terminal initiation site, or vice versa.

As demonstrated by the examples described herein, recombinant RSV genome in the positive-sense or negative-sense orientation is co-transfected with expression vectors encoding the viral nucleocapsid (N) protein, the associated nucleocapsid phosphoprotein (P), the large (L) polymerase subunit protein, with or without the M2/ORF1 protein of RSV to generate infectious viral particles. Vaccinia vectors expressing RSV virus polypeptides are used as the source of proteins which were able to replicate and transcribe synthetically derived RNPs. The minimum subset of RSV proteins needed for specific replication and expression of the viral RNP was found to be the three polymerase complex proteins (N, P and L). This suggests that the M2 gene function is not absolutely required for the replication, expression and rescue of infectious RSV.

The expression products and/or chimeric virions obtained may advantageously be utilized in vaccine formulations. In particular, recombinant RSV genetically engineered to demonstrate an attenuated phenotype may be utilized as a live



RSV vaccine. In another embodiment of the invention, recombinant RSV may be engineered to express the antigenic polypeptides of another strain of RSV (e.g., RSV G and F proteins) or another virus (e.g., an immunogenic peptide from 5 gp120 of HIV) to generate a chimeric RSV to serve as a vaccine, that is able to elicit both vertebrate humoral and cell-mediated immune responses. The use of recombinant influenza or recombinant RSV for this purpose is especially attractive since these viruses demonstrate tremendous strain 10 variability allowing for the construction of a vast repertoire of vaccine formulations. The ability to select from thousands of virus variants for constructing chimeric viruses obviates the problem of host resistance encountered when using other viruses such as vaccinia.

15

### 3.1. DEFINITIONS

As used herein, the following terms will have the meanings indicated:

- crNA = anti-genomic RNA
- 20 HA = hemagglutinin (envelope glycoprotein)
- HIV = human immunodeficiency virus
- L = large polymerase subunit
- M = matrix protein (lines inside of envelope)
- MDCK = Madin Darby canine kidney cells
- 25 MDBK = Madin Darby bovine kidney cells
- moi = multiplicity of infection
- N = nucleocapsid protein
- NA = neuraminidase (envelope glycoprotein)
- NP = nucleoprotein (associated with RNA and required 30 for polymerase activity)
- NS = nonstructural protein (function unknown)
- nt = nucleotide
- P = nucleocapsid phosphoprotein
- PA, PB1, PB2 = RNA-directed RNA polymerase components
- 35 RNP = ribonucleoprotein (RNA, PB2, PB1, PA and NP)
- rRNP = recombinant RNP
- RSV = respiratory syncytial virus

vRNA = genomic virus RNA  
 viral polymerase complex = PA, PB1, PB2 and NP  
 WSN = influenza A/WSN/33 virus  
 WSN-HK virus: reassortment virus containing seven genes  
 from WSN virus and the NA gene from  
 influenza A/HK/8/68 virus

#### 4. DESCRIPTION OF THE FIGURES

FIG. 1. Schematic representation of the RSV/CAT  
 construct (pRSVA2CAT) used in rescue experiments. The  
 approximate 100 nt long leader and 200 nt long trailer regions  
 of RSV were constructed by the controlled annealing of  
 synthetic oligonucleotides containing partial overlapping  
 complementarity. The overlapping leader oligonucleotides are  
 indicated by the 1L - 5L shown in the construct. The  
 overlapping trailer nucleotides are indicated by the 1T - 9T  
 shown in the construct. The nucleotide sequences of the  
 leader and trailer DNAs were ligated into purified CAT gene  
 DNA at the indicate XbaI and PstI sites respectively. This  
 entire construct was then ligated into KpnI/HindIII digested  
 pUC19. The inclusion of a T7 promoter sequence and a HgaI  
 site flanking the trailer and leader sequences, respectively,  
 allowed in vitro synthesis of RSV/CAT RNA transcripts  
 containing the precise genomic sequence 3' and 5' ends.

FIG. 2. Thin layer chromatogram (TLC) showing the CAT  
 activity present in 293 cell extracts following infection and  
 transfection with RNA transcribed from the RSV/CAT construct  
 shown in Figure 1. Confluent monolayers of 293 cells in six-  
 well plates ( $\sim 10^6$  cells) were infected with either RSV A2 or  
 B9320 at an m.o.i. of 0.1-1.0 pfu cell. At 1 hour post  
 infection cells were transfected with 5-10  $\mu$ g of CAT/RSV using  
 the Transfect-Act<sup>™</sup> protocol of Life Technologies. At 24 hours  
 post infection the infected/transfected monolayers were  
 harvested and processed for subsequence CAT assay according to  
 Current Protocols in Molecular Biology, Vol. 1, Chapter 9.6.2;  
 Gorman, et al., (1982) Mol. Cell. Biol. 2:1044-1051. Lanes 1,

2, 3 and 4 show the CAT activity present in (1) uninfected 293 cells, transfected with CAT/RSV-A2 infected 293 cells, co-infected with supernatant from (2) above. The CAT activity observed in each lane was produced from 1/5 of the total cellular extract from  $10^6$  cells.

FIG. 3. Schematic representation of the RSV strain A2 genome showing the relative positions of the primer pairs used for the synthesis of cDNAs comprising the entire genome. The endonuclease sites used to splice these clones together are indicated; these sites were present in the native RSV sequence and were included in the primers used for cDNA synthesis. Approximately 100 ng of viral genomic RNA was used in RT/PCR reactions for the separate synthesis of each of the seven cDNAs. The primers for the first and second strand cDNA synthesis from the genomic RNA template are also shown. For each cDNA, the primers for the first strand synthesis are nos. 1-7 and the primers for the second strand synthesis are nos. 1'-7'.

20

FIG. 4. Schematic representation of the RSV subgroup B strain B9320. BamHI sites were created in the oligonucleotide primers used for RT/PCR in order to clone the G and F genes from the B9320 strain into RSV subgroup A2 antigenomic cDNA (FIG. 4A). A cDNA fragment which contained G and F genes from 4326 nucleotides to 9387 nucleotides of A2 strain was first subcloned into pUC19 (pUCRVH). Bgl II sites were created at positions of 4630 (SH/G intergenic junction FIG. 4B) and 7554 (F/M2 intergenic junction (FIG. 4C). B93260 A-G and -F cDNA inserted into pUCR/H which is deleted of the A-G and F genes. The resulting antigenomic cDNA clone was termed as pRSVB-GF and was used to transfect Hep-2 cells to generate infectious RSVB-GF virus.

FIG. 5. Recombinant RSVB-GF virus was characterized by RT/PCR using RSV subgroup B specific primers. RSV subgroup B specific primers in the G region were incubated with aliquots

of the recombinant RSV viral genomes and subjected to PCR. The PCR products were analyzed by electrophoresis on a 1% agarose gel and visualized by staining with ethidium bromide. As shown, no DNA product was produced in the RT/PCR reaction using RSV A2 as a template. However, a predicted product of 254 base pairs was seen in RT/PCR of RSVB-GF RNA and PCR control of plasmid pRSV-GF DNA as template, indicating the rescued virus contained G and F genes derived from B9320 virus.

10

FIG. 6. Identification of chimeric (rRSVA2(B-G) by RT/PCR and Northern blot analysis of RNA expression. FIG. 6A. RT/PCR analysis of chimeric rRSV A2(B-G), A2(B-G), in comparison with wild-type A2(A2). Virion RNA extracted from 15 rRSVA2(B-G) (lanes 1, 2) and rRSVA2 (lanes 3,4) was reverse transcribed using a primer annealed to (-) sense vRNA in the RSV F gene in the presence (+) or absence (-) of reverse transcriptase (RT), followed by PCR with a primer pair flanking the B-G insertion site. No DNA was detected in 20 RT/PCR when reverse transcriptase (RT) was absent (lanes 2,4). A cDNA fragment, which is about 1kb bigger than the cDNA derived from A2, was produced from rRSVA(B-G). This longer PCR DNA product was digested by *Stu I* restriction enzyme unique to the inserted B-G gene (lane 5). 100 bp DNA size 25 marker is indicated (M). FIG. 6B. Northern blot analysis of G mRNA expression. Hep-2 cells were infected with RSV B9320, rRSV and chimeric rRSV A2 (B-G). At 48 hr postinfection, total cellular RNA was extracted and electrophoresed on a 1.2% agarose gel containing formaldehyde. RNA was transferred to 30 Hybond Nylon membrane and the filter was hybridized with a <sup>32</sup>P-labeled oligonucleotide probe specific for A2-G or specific for B9320-G mRNA. Both A2 G specific and B9320 G specific transcripts were detected in the rRSV A2 (B-G) infected cells. The run-off RNA transcript (G-M2) from rRSV A2 35 (B-G) infected cells is also indicated.

FIG. 7. Analysis of protein expression by rRSV A2 (B-G). Hep-2 cells were mock-infected (lanes 1, 5), infected with RSV B9320 (lanes 2, 6), rRSV (lanes 3, 7) and rRSV A2 (B-G) (lanes 4, 8). At 14-18 hr postinfection, infected cells were labeled with  $^{35}\text{S}$ -promix and polypeptides were immunoprecipitated by goat polyclonal antiserum against RSV A2 strain (lanes 1-5) or by mouse polyclonal antiserum against RSV B9320 strain (lanes 5-8). Immunoprecipitated polypeptides were separated on a 10% polyacrylamide gel. Both RSV A2 specific G protein and RSV B9320 specific G protein were produced in rRSV A2 (B-G) infected cells. The G protein migration is indicated by \*. Mobility of the F1 glycoprotein, and N, P, and M is indicated. Molecular sizes are shown on the left in kilodaltons.

FIG. 8. Plaque morphology of rRSV, rRSVC3G, rRSV A2 (B-G) and wild-type A2 virus (wt A2). Hep-2 cells were infected with each virus and incubated at 35°C for six days. The cell monolayers were fixed, visualized by immunostaining, and photographed.

FIG. 9. Growth curve of rRSV, rRSVC4G, wild-type A2 RSV (wt A2) and chimeric rRSV A2 (B-G). Hep-2 cells were infected with either virus at a moi of 0.5 and the medium was harvested at 24 hr intervals. The titer of each virus was determined in duplicate by plaque assay on Hep-2 cells and visualized by immunostaining.

FIG. 10. RSV L protein charged residue clusters targeted for site-directed mutagenesis. Charged amino acid residues in contiguous clusters were converted to alanines by site-directed mutagenesis of the RSV L gene using the QuikChange site-directed mutagenesis kit (Stratagene).

FIG. 11. RSV L protein cysteine residues targeted for site-directed mutagenesis. Cysteine residues were converted to alanine-residues by site-directed mutagenesis of the RSV L

gene using the QuikChange site-directed mutagenesis kit (Stratagene).

FIG. 12. Identification RSV M2-2 and SH deletion mutants. Deletions in M2-2 were generated by Hind III digestion of pET(S/B) followed by recloning of a remaining Sac I to BamHI fragment into a full-length clone. Deletions in SH were generated by Sac I digestion of pET(A/S) followed by recloning of a remaining Avr II Sac I fragment into a full-length clone. FIG. 12A. Identification of the recovered rRSVsSH and rRSV M2-2 was performed by RT/PCR using primer pairs specific for the SH gene or M2-2 gene, respectively. Fig. 12B rRSV SH M2-2 was also detected by RT/PCR using primer pairs specific for the M2-2 and SH genes. RT/PCR products were run on an ethidium bromide agarose gel and bands were visualized by ultraviolet (UV) light.

#### 5. DESCRIPTION OF THE INVENTION

The present invention relates to genetically engineered recombinant RS viruses and viral vectors which express heterologous genes or mutated RS viral genes or a combination of viral genes derived from different strains of RS virus. The invention relates to the construction and use of recombinant negative strand RS viral RNA templates which may be used with viral RNA-directed RNA polymerase to express heterologous gene products in appropriate host cells and/or to rescue the heterologous gene in virus particles. The RNA templates of the present invention may be prepared by transcription of appropriate DNA sequences using a DNA-directed RNA polymerase such as bacteriophage T7, T3 or Sp6 polymerase. The recombinant RNA templates may be used to transfect continuous/transfected cell lines that express the RNA-directed RNA polymerase proteins allowing for complementation.

The invention is demonstrated by way of working examples in which infectious RSV is rescued from cDNA containing the

RSV genome in the genomic or antigenomic sense introduced into cells expressing the N, P, and L proteins of the RSV polymerase complex. The working examples further demonstrate that expression of M2-1 is not required for recovery of  
5 infectious RSV from cDNA which is contrary to what has been reported earlier (Collins et al., 1995, Proc. Natl. Acad. Sci. USA 92:11563-7). Furthermore, the addition of plasmids expressing M2-1 has little effect on the RSV rescue efficiency. M2-deleted-RSV is an excellent vehicle to  
10 generate chimeric RSV encoding heterologous gene products in place of the M2 genes, these chimeric viral vectors and rescued virus particles have utility as expression vectors for the expression of heterologous gene products and as live attenuated RSV vaccines expressing either RSV antigenic  
15 polypeptides or antigenic polypeptides of other viruses.

The invention is further demonstrated by way of working examples in which a cDNA clone which contained the complete genome of RSV, in addition to a T7 promoter, a hepatitis delta virus ribozyme and a T7 terminator is used to generate an  
20 infectious viral particle when co-transfected with expression vectors encoding the N, P, L and M2/orf1 proteins of RSV. In addition, the working examples describe RNA transcripts of cloned DNA containing the coding region -- in negative sense orientation -- of the chloramphenicol-acetyl-transferase (CAT)  
25 gene or the green fluorescent protein (GFP) gene flanked by the 5' terminal and 3' terminal nucleotides of the RSV genome. The working examples further demonstrate that an RSV promoter mutated to have increased activity resulted in rescue of infectious RSV particles from a full length RSV cDNA with high  
30 efficiency. These results demonstrate the successful use of recombinant viral negative strand templates and RSV polymerase with increased activity to rescue RSV. This system is an excellent tool to engineer RSV viruses with defined biological properties, e.g. live-attenuated vaccines against RSV, and to  
35 use recombinant RSV as an expression vector for the expression of heterologous gene products.

This invention relates to the construction and use of recombinant negative strand viral RNA templates which may be used with viral RNA-directed RNA polymerase to express heterologous gene products in appropriate host cells, to  
5 rescue the heterologous gene in virus particles and/or express mutated or chimeric recombinant negative strand viral RNA templates (see U.S. Patent No. 5,166,057 to Palese et al., incorporated herein by reference in its entirety). In a specific embodiment of the invention, the heterologous gene  
10 product is a peptide or protein derived from another strain of the virus or another virus. The RNA templates may be in the positive or negative-sense orientation and are prepared by transcription of appropriate DNA sequences using a DNA-directed RNA polymerase such as bacteriophage T7, T3 or the  
15 Sp6 polymerase.

The ability to reconstitute RNP's in vitro allows the design of novel chimeric influenza and RSV viruses which express foreign genes. One way to achieve this goal involves modifying existing viral genes. For example, the HA gene of  
20 influenza may be modified to contain foreign sequences in its external domains. Where the heterologous sequence are epitopes or antigens of pathogens, these chimeric viruses may be used to induce a protective immune response against the disease agent from which these determinants are derived.  
25 For example, a chimeric RNA may be constructed in which a coding sequence derived from the gp120 coding region of human immunodeficiency virus was inserted into the F or G coding sequence of influenza, and chimeric virus was produced from transfection of this chimeric RNA segment into a host cell  
30 infected with wild-type RSV.

In addition to modifying genes coding for surface proteins, genes coding for nonsurface proteins may be altered. The latter genes have been shown to be associated with most of the important cellular immune responses in the RS virus  
35 system. Thus, the inclusion of a foreign determinant in the G or F gene of RSV may - following infection - induce an effective cellular immune response against this determinant.



Such an approach may be particularly helpful in situations in which protective immunity heavily depends on the induction of cellular immune responses (e.g., malaria, etc.).

- The present invention also relates to attenuated
- 5 recombinant RSV produced by introducing specific mutations in the genome of RSV which results in an amino acid change in an RSV protein, such as a polymerase protein, which results in an attenuated phenotype.

10           **5.1. CONSTRUCTION OF THE RECOMBINANT RNA TEMPLATES**

- Heterologous gene coding sequences flanked by the complement of the viral polymerase binding site/promoter, e.g., the complement of the 3'-RSV termini or the 3'- and 5'- RSV termini may be constructed using techniques known in the art.
- 15 Heterologous gene coding sequences may also be flanked by the complement of the RSV polymerase binding site/promoter, e.g., the leader and trailer sequence of RSV using techniques known in the art. Recombinant DNA molecules containing these hybrid sequences can be cloned and transcribed by a DNA-directed RNA
- 20 polymerase, such as bacteriophage T7, T3 or the Sp6 polymerase and the like, to produce the recombinant RNA templates which possess the appropriate viral sequences that allow for viral polymerase recognition and activity.

- In a preferred embodiment of the present invention, the
- 25 heterologous sequences are derived from the genome of another strain of RSV, e.g., the genome of RSV A strain is engineered to include the nucleotide sequences encoding the antigenic polypeptides G and F of RSV B strain, or fragments thereof. In such an embodiment of the invention, heterologous coding
- 30 sequences from another strain of RSV can be used to substitute for nucleotide sequences encoding antigenic polypeptides of the starting strain, or be expressed in addition to the antigenic polypeptides of the parent strain, so that a recombinant RSV genome is engineered to express the antigenic
- 35 polypeptides of one, two or more strains of RSV.

In yet another embodiment of the invention, the heterologous sequences are derived from the genome of any

strain of influenza virus. In accordance with the present invention, the heterologous coding sequences of influenza may be inserted within a RSV coding sequence such that a chimeric gene product is expressed which contains the heterologous peptide sequence within the RSV viral protein. In either embodiment, the heterologous sequences derived from the genome of influenza may include, but are not limited to HA, NA, PB1, PB2, PA, NS1 or NS2.

- In one specific embodiment of the invention, the heterologous sequences are derived from the genome of human immunodeficiency virus (HIV), preferably human immunodeficiency virus-1 or human immunodeficiency virus-2. In another embodiment of the invention, the heterologous coding sequences may be inserted within an influenza gene coding sequence such that a chimeric gene product is expressed which contains the heterologous peptide sequence within the influenza viral protein. In such an embodiment of the invention, the heterologous sequences may also be derived from the genome of a human immunodeficiency virus, preferably of human immunodeficiency virus-1 or human immunodeficiency virus-2.

- In instances whereby the heterologous sequences are HIV-derived, such sequences may include, but are not limited to sequences derived from the env gene (i.e., sequences encoding all or part of gp160, gp120, and/or gp41), the pol gene (i.e., sequences encoding all or part of reverse transcriptase, endonuclease, protease, and/or integrase), the gag gene (i.e., sequences encoding all or part of p7, p6, p55, p17/18, p24/25) tat, rev, nef, vif, vpu, vpr, and/or vpx.

- One approach for constructing these hybrid molecules is to insert the heterologous coding sequence into a DNA complement of a RSV genomic RNA so that the heterologous sequence is flanked by the viral sequences required for viral polymerase activity; i.e., the viral polymerase binding site/promoter, hereinafter referred to as the viral polymerase binding site. In an alternative approach, oligonucleotides encoding the viral polymerase binding site, e.g., the

complement of the 3'-terminus or both termini of the virus genomic segments can be ligated to the heterologous coding sequence to construct the hybrid molecule. The placement of a foreign gene or segment of a foreign gene within a target  
5 sequence was formerly dictated by the presence of appropriate restriction enzyme sites within the target sequence. However, recent advances in molecular biology have lessened this problem greatly. Restriction enzyme sites can readily be placed anywhere within a target sequence through the use of  
10 site-directed mutagenesis (e.g., see, for example, the techniques described by Kunkel, 1985, Proc. Natl. Acad. Sci. U.S.A. 82;488). Variations in polymerase chain reaction (PCR) technology, described infra, also allow for the specific insertion of sequences (i.e., restriction enzyme sites) and  
15 allow for the facile construction of hybrid molecules. Alternatively, PCR reactions could be used to prepare recombinant templates without the need of cloning. For example, PCR reactions could be used to prepare double-stranded DNA molecules containing a DNA-directed RNA  
20 polymerase promoter (e.g., bacteriophage T3, T7 or Sp6) and the hybrid sequence containing the heterologous gene and the influenza viral polymerase binding site. RNA templates could then be transcribed directly from this recombinant DNA. In yet another embodiment, the recombinant RNA templates may be  
25 prepared by ligating RNAs specifying the negative polarity of the heterologous gene and the viral polymerase binding site using an RNA ligase. Sequence requirements for viral polymerase activity and constructs which may be used in accordance with the invention are described in the subsections  
30 below.

#### 5.1.1. INSERTION OF THE HETEROLOGOUS GENES

The genes coding for the M2 or L proteins contain a single open reading frame. The gene coding for NS contains  
35 two open reading frames for NS1 and NS2. The G and F proteins, coded for by separate genes, are the major surface glycoproteins of the virus. Consequently, these proteins are

the major targets for the humoral immune response after infection. Insertion of a foreign gene sequence into any of these coding regions could be accomplished by either a complete replacement of the viral coding region with the  
5 foreign gene or by a partial replacement. Complete replacement would probably best be accomplished through the use of PCR-directed mutagenesis.

Alternatively, a bicistronic mRNA could be constructed to permit internal initiation of translation of viral sequences  
10 and allow for the expression of foreign protein coding sequences from the regular terminal initiation site. Alternatively, a bicistronic mRNA sequence may be constructed wherein the viral sequence is translated from the regular terminal open reading frame, while the foreign sequence is  
15 initiated from an internal site. Certain internal ribosome entry site (IRES) sequences may be utilized. The IRES sequences which are chosen should be short enough to not interfere with RS virus packaging limitations. Thus, it is preferable that the IRES chosen for such a bicistronic  
20 approach be no more than 500 nucleotides in length, with less than 250 nucleotides being preferred. Further, it is preferable that the IRES utilized not share sequence or structural homology with picornaviral elements. Preferred IRES elements include, but are not limited to the mammalian  
25 BiP IRES and the hepatitis C virus IRES.

## 5.2. EXPRESSION OF HETEROLOGOUS GENE PRODUCTS USING RECOMBINANT RNA TEMPLATE

The recombinant templates prepared as described above can  
30 be used in a variety of ways to express the heterologous gene products in appropriate host cells or to create chimeric viruses that express the heterologous gene products. In one embodiment, the recombinant template can be combined with viral polymerase complex purified *infra*, to produce rRNPs  
35 which are infectious. To this end, the recombinant template can be transcribed in the presence of the viral polymerase complex. Alternatively, the recombinant template may be mixed

with or transcribed in the presence of viral polymerase complex prepared using recombinant DNA methods (e.g. see Kingsbury et al., 1987, Virology 156:396-403). In yet another embodiment, the recombinant template can be used to transfect appropriate host cells to direct the expression of the heterologous gene product at high levels. Host cell systems which provide for high levels of expression include continuous cell lines that supply viral functions such as cell lines superinfected with RSV, cell lines engineered to complement RSV viral functions, etc.

### 5.3. PREPARATION OF CHIMERIC NEGATIVE STRAND RNA VIRUS

In order to prepare chimeric virus, reconstituted RNPs containing modified RSV RNAs or RNA coding for foreign proteins may be used to transfect cells which are also infected with a "parent" RSV virus. Alternatively, the reconstituted RNP preparations may be mixed with the RNPs of wild type parent virus and used for transfection directly. Following reassortment, the novel viruses may be isolated and their genomes be identified through hybridization analysis. In additional approaches described herein for the production of infectious chimeric virus, rRNPs may be replicated in host cell systems that express the RSV or influenza viral polymerase proteins (e.g., in virus/host cell expression systems; transformed cell lines engineered to express the polymerase proteins, etc.), so that infectious chimeric virus are rescued; in this instance, helper virus need not be utilized since this function is provided by the viral polymerase proteins expressed. In a particularly desirable approach, cells infected with rRNPs engineered for all eight influenza virus segments may result in the production of infectious chimeric virus which contain the desired genotype; thus eliminating the need for a selection system.

Theoretically, one can replace any one of the genes of RSV, or part of any one of the RSV genes, with the foreign sequence. However, a necessary part of this equation is the

ability to propagate the defective virus (defective because a normal viral gene product is missing or altered). A number of possible approaches exist to circumvent this problem.

- A third approach to propagating the recombinant virus may
- 5 involve co-cultivation with wild-type virus. This could be done by simply taking recombinant virus and co-infecting cells with this and another wild-type virus (preferably a vaccine strain). The wild-type virus should complement for the defective virus gene product and allow growth of both the
- 10 wild-type and recombinant virus. This would be an analogous situation to the propagation of defective-interfering particles of influenza virus (Nayak et al., 1983, In: Genetics of Influenza Viruses, P. Palese and D. W. Kingsbury, eds., Springer-Verlag, Vienna, pp. 255-279). In the case of
- 15 defective-interfering viruses, conditions can be modified such that the majority of the propagated virus is the defective particle rather than the wild-type virus. Therefore this approach may be useful in generating high titer stocks of recombinant virus. However, these stocks would necessarily
- 20 contain some wild-type virus.

- Alternatively, synthetic RNPs may be replicated in cells co-infected with recombinant viruses that express the RS virus polymerase proteins. In fact, this method may be used to rescue recombinant infectious virus in accordance with the
- 25 invention. To this end, the RSV virus polymerase proteins may be expressed in any expression vector/host cell system, including but not limited to viral expression vectors (e.g., vaccinia virus, adenovirus, baculovirus, etc.) or cell lines that express the polymerase proteins (e.g., see Krystal et
- 30 al., 1986, Proc. Natl. Acad. Sci. USA 83: 2709-2713).

#### 5.4. GENERATION OF CHIMERIC VIRUSES WITH AN ATTENUATED PHENOTYPE

- The methods of present invention may be used to introduce
- 35 mutations or heterologous sequences to generate chimeric attenuated viruses which have many applications, including analysis of RSV molecular biology, pathogenesis, and growth

and infection properties. In accordance with the present invention, mutations or heterologous sequences may be introduced for example into the F or G protein coding sequences, NS1, NS2, M1ORF, M2ORF, N, P, or L coding sequences, NS1, NS2, M1ORF, M2ORF, N, P, or L coding

5 sequences. In yet another embodiment of the present invention, a particular viral gene, or the expression thereof, may be eliminated to generate an attenuated phenotype, e.g., the M ORF may be deleted from the RSV genome to generate a recombinant RSV with an attenuated phenotype. In yet another

10 embodiment, the individual internal genes of human RSV can be replaced by another strains counterpart, or their bovine or murine counterpart. This may include part or all of one or more of the NS1, NS2, N, P, M, SH, M2(ORF1), M2(ORF2) and L genes or the G and F genes. The RSV genome contains ten mRNAs

15 encoding three transmembrane proteins, G protein, fusion F protein required for penetration, and the small SH protein; the nucleocapsid proteins N, P and L; transcription elongation factor M2 ORF 1; the matrix M protein and two nonstructural proteins, NS1 and NS2. Any one of the proteins may be

20 targeted to generate and attenuated phenotype. Other mutations which may be utilized to result in an attenuated phenotype are insertional, deletional and site directed mutations of the leader and trailer sequences.

In accordance with the present invention, an attenuated

25 RSV exhibits a substantially lower degree of virulence as compared to a wild-type virus, including a slower growth rate, such that the symptoms of viral infection do not occur in an immunized individual.

In accordance with the present invention attenuated

30 recombinant RSV may be generated by incorporating a broad range of mutations including single nucleotide changes, site-specific mutations, insertions, substitutions, deletions, or rearrangements. These mutations may affect a small segment of the RSV genome, e.g., 15 to 30 nucleotides, or large segments

35 of the RSV genome, e.g., 50 to 1000 nucleotides, depending on the nature of the mutation. In yet another embodiment, mutations are introduced upstream or downstream of an existing

cis-acting regulatory element in order to ablate its activity, thus resulting in an attenuated phenotype.

In accordance with the invention, a non-coding regulatory region of a virus can be altered to down-regulate any viral gene, e.g. reduce transcription of its mRNA and/or reduce replication of vRNA (viral RNA), so that an attenuated virus is produced.

Alterations of non-coding regulatory regions of the viral genome which result in down-regulation of replication of a viral gene, and/or down-regulation of transcription of a viral gene will result in the production of defective particles in each round of replication; i.e. particles which package less than the full complement of viral segments required for a fully infectious, pathogenic virus. Therefore, the altered virus will demonstrate attenuated characteristics in that the virus will shed more defective particles than wild type particles in each round of replication. However, since the amount of protein synthesized in each round is similar for both wild type virus and the defective particles, such attenuated viruses are capable of inducing a good immune response.

The foregoing approach is equally applicable to both segmented and non-segmented viruses, where the down regulation of transcription of a viral gene will reduce the production of its mRNA and the encoded gene product. Where the viral gene encodes a structural protein, e.g., a capsid, matrix, surface or envelope protein, the number of particles produced during replication will be reduced so that the altered virus demonstrates attenuated characteristics; e.g., a titer which results in subclinical levels of infection. For example, a decrease in viral capsid expression will reduce the number of nucleocapsids packaged during replication, whereas a decrease in expression of the envelope protein may reduce the number and/or infectivity of progeny virions. Alternatively, a decrease in expression of the viral enzymes required for replication, e.g., the polymerase, replicase, helicase, and the like, should decrease the number of progeny genomes



generated during replication. Since the number of infectious particles produced during replication are reduced, the altered viruses demonstrated attenuated characteristics. However, the number of antigenic virus particles produced will be

5 sufficient to induce a vigorous immune response.

An alternative way to engineer attenuated viruses involves the introduction of an alteration, including but not limited to an insertion, deletion or substitution of one or more amino acid residues and/or epitopes into one or more of  
10 the viral proteins. This may be readily accomplished by engineering the appropriate alteration into the corresponding viral gene sequence. Any change that alters the activity of the viral protein so that viral replication is modified or reduced may be accomplished in accordance with the invention.

15 For example, alterations that interfere with but do not completely abolish viral attachment to host cell receptors and ensuing infection can be engineered into viral surface antigens or viral proteases involved in processing to produce an attenuated strain. According to this embodiment, viral  
20 surface antigens can be modified to contain insertions, substitution or deletions of one or more amino acids or epitopes that interfere with or reduce the binding affinity of the viral antigen for the host cell receptors. This approach offers an added advantage in that a chimeric virus which  
25 expresses a foreign epitope may be produced which also demonstrates attenuated characteristics. Such viruses are ideal candidates for use as live recombinant vaccines. For example, heterologous gene sequences that can be engineered into the chimeric viruses of the invention include, but are  
30 not limited to, epitopes of human immunodeficiency virus (HIV) such as gp120; hepatitis B virus surface antigen (HBsAg); the glycoproteins of herpes virus (e.g., gD, gE); VP1 of poliovirus; and antigenic determinants of nonviral pathogens such as bacteria and parasites, to name but a few.

35 In this regard, RSV is an ideal system in which to engineer foreign epitopes, because the ability to select from thousands of virus variants for constructing chimeric viruses

obviates the problem of host resistance or immune tolerance encountered when using other virus vectors such as vaccinia.

In another embodiment, alterations of viral proteases required for processing viral proteins can be engineered to  
5 produce attenuation. Alterations which affect enzyme activity and render the enzyme less efficient in processing, should affect viral infectivity, packaging, and/or release to produce an attenuated virus.

In another embodiment, viral enzymes involved in viral  
10 replication and transcription of viral genes, e.g., viral polymerases, replicases, helicases, etc. may be altered so that the enzyme is less efficient or active. Reduction in such enzyme activity may result in the production of fewer progeny genomes and/or viral transcripts so that fewer  
15 infectious particles are produced during replication.

The alterations engineered into any of the viral enzymes include but are not limited to insertions, deletions and substitutions in the amino acid sequence of the active site of the molecule. For example, the binding site of the enzyme  
20 could be altered so that its binding affinity for substrate is reduced, and as a result, the enzyme is less specific and/or efficient. For example, a target of choice is the viral polymerase complex since temperature sensitive mutations exist in all polymerase proteins. Thus, changes introduced into the  
25 amino acid positions associated with such temperature sensitivity can be engineered into the viral polymerase gene so that an attenuated strain is produced.

#### 5.4.1. THE RSV L GENE AS A TARGET FOR ATTENUATION

30 In accordance with the present invention, the RSV L gene is an important target to generate recombinant RSV with an attenuated phenotype. The L gene represents 48% of the entire RSV genome. The present invention encompasses generating L  
35 gene mutants with defined mutations or random mutations in the RSV L gene. Any number of techniques known to those skilled in the art may be used to generate both defined or random

mutations into the RSV L gene. Once the mutations have been introduced, the functionality of the L gene cDNA mutants are screened in vitro using a minigenome replication system and the recovered L gene mutants are then further analyzed in

5 vitro and in vivo.

The following strategies are exemplary of the approaches which may be used to generate mutants with an attenuated phenotype. Further, the following strategies as described below have been applied to the L gene only by way of example  
10 and may also be applied to any of the other RSV genes.

One approach to generate mutants with an attenuated phenotype utilizes a scanning mutagenesis approach to mutate clusters of charged amino acids to alanines. This approach is particularly effective in targeting functional domains, since  
15 the clusters of charged amino acids generally are not found buried within the protein structure. Replacing the charged amino acids with conservative substitutions, such as neutral amino acids, e.g., alanine, should not grossly alter the structure of the protein but rather, should alter the activity  
20 of the functional domain of the protein. Thus, disruption of charged clusters should interfere with the ability of that protein to interact with other proteins, thus making the mutated protein's activity thermosensitive which can yield temperature sensitive mutants.

25 A cluster of charged amino acids may be arbitrarily defined as a stretch of five amino acids in which at least two or more residues are charged residues. In accordance with the scanning mutagenesis approach all of the charged residues in the cluster are mutated to alanines using site-directed  
30 mutagenesis. Due to the large size of the RSV L gene, there are many clustered charged residues. Within the L gene, there are at least two clusters of four contiguous charged residues and at least seventeen clusters of three contiguous charged residues. At least two to four of the charged residues in  
35 each cluster may be substituted with a neutral amino acid, e.g., alanine.

In yet another approach to generate mutants with an attenuated phenotype utilizes a scanning mutagenesis approach to mutate cysteines to amino acids, such as glycines or alanines. Such an approach takes advantage of the frequent  
5 role of cysteines in intramolecular and intermolecular bond formations, thus by mutating cysteines to another residue, such as a conservative substitution e.g., valine or alanine, or a drastic substitution e.g., aspartic acid, the stability and function of a protein may be altered due to disruption of  
10 the protein's tertiary structure. There are approximately thirty-nine cysteine residues present in the RSV L gene.

In yet another approach random mutagenesis of the RSV L gene will cover residues other than charged or cysteines. Since the RSV L gene is very large, such an approach may be  
15 accomplished by mutagenizing large cDNA fragments of the L gene by PCR mutagenesis. The functionality of such mutants may be screened by a minigenome replication system and the recovered mutants are then further analyzed in vitro and in vivo.

20

#### 5.5. VACCINE FORMULATIONS USING THE CHIMERIC VIRUSES

Virtually any heterologous gene sequence may be constructed into the chimeric viruses of the invention for use  
25 in vaccines. In a preferred embodiment, the present invention relates to bivalent RSV vaccines which confers protection against RSV-A and RSV-B. To formulate such a vaccine, a chimeric RS virus is used which expresses the antigenic polypeptides of both RSV-A and RSV-B subtypes. In yet another  
30 preferred embodiment, the present invention relates to a bivalent vaccine which confers protection against both RSV and influenza. To formulate such a vaccine, a chimeric RS virus is used which expresses the antigenic polypeptides of both RSV and influenza.

35 Preferably, epitopes that induce a protective immune response to any of a variety of pathogens, or antigens that bind neutralizing antibodies may be expressed by or as part of

chimeric viruses. For example, heterologous gene sequences that can be constructed into the chimeric viruses of the invention for use in vaccines include but are not limited to sequences derived from a human immunodeficiency virus (HIV),  
5 preferably type 1 or type 2. In a preferred embodiment, an immunogenic HIV-derived peptide which may be the source of an antigen may be constructed into a chimeric influenza virus that may then be used to elicit a vertebrate immune response.

Such HIV-derived peptides may include, but are not  
10 limited to sequences derived from the env gene (i.e., sequences encoding all or part of gp160, gp120, and/or gp41), the pol gene (i.e., sequences encoding all or part of reverse transcriptase, endonuclease, protease, and/or integrase), the gag gene (i.e., sequences encoding all or part of p7, p6, p55,  
15 p17/18, p24/25), tat, rev, nef, vif, vpr, and/or vpx.

Other heterologous sequences may be derived from hepatitis B virus surface antigen (HBsAg); the glycoproteins of herpes virus (e.g. gD, gE); VP1 of poliovirus; antigenic determinants of non-viral pathogens such as bacteria and  
20 parasites, to name but a few. In another embodiment, all or portions of immunoglobulin genes may be expressed. For example, variable regions of anti-idiotypic immunoglobulins that mimic such epitopes may be constructed into the chimeric viruses of the invention.

25 Either a live recombinant viral vaccine or an inactivated recombinant viral vaccine can be formulated. A live vaccine may be preferred because multiplication in the host leads to a prolonged stimulus of similar kind and magnitude to that occurring in natural infections, and therefore, confers  
30 substantial, long-lasting immunity. Production of such live recombinant virus vaccine formulations may be accomplished using conventional methods involving propagation of the virus in cell culture or in the allantois of the chick embryo followed by purification.

35 In this regard, the use of genetically engineered RSV (vectors) for vaccine purposes may require the presence of attenuation characteristics in these strains. Current live

virus vaccine candidates for use in humans are either cold adapted, temperature sensitive, or passaged so that they derive several (six) genes from avian viruses, which results in attenuation. The introduction of appropriate mutations 5 (e.g., deletions) into the templates used for transfection may provide the novel viruses with attenuation characteristics. For example, specific missense mutations which are associated with temperature sensitivity or cold adaption can be made into deletion mutations. These mutations should be more stable 10 than the point mutations associated with cold or temperature sensitive mutants and reversion frequencies should be extremely low.

Alternatively, chimeric viruses with "suicide" characteristics may be constructed. Such viruses would go 15 through only one or a few rounds of replication in the host. When used as a vaccine, the recombinant virus would go through a single replication cycle and induce a sufficient level of immune response but it would not go further in the human host and cause disease. Recombinant viruses lacking one or more of 20 the essential RSV virus genes would not be able to undergo successive rounds of replication. Such defective viruses can be produced by co-transfecting reconstituted RNPs lacking a specific gene(s) into cell lines which permanently express this gene(s). Viruses lacking an essential gene(s) will be 25 replicated in these cell lines but when administered to the human host will not be able to complete a round of replication. Such preparations may transcribe and translate -in this abortive cycle -- a sufficient number of genes to induce an immune response. Alternatively, larger quantities 30 of the strains could be administered, so that these preparations serve as inactivated (killed) virus vaccines. For inactivated vaccines, it is preferred that the heterologous gene product be expressed as a viral component, so that the gene product is associated with the virion. The 35 advantage of such preparations is that they contain native proteins and do not undergo inactivation by treatment with

formalin or other agents used in the manufacturing of killed virus vaccines.

In another embodiment of this aspect of the invention, inactivated vaccine formulations may be prepared using 5 conventional techniques to "kill" the chimeric viruses. Inactivated vaccines are "dead" in the sense that their infectivity has been destroyed. Ideally, the infectivity of the virus is destroyed without affecting its immunogenicity. In order to prepare inactivated vaccines, the chimeric virus 10 may be grown in cell culture or in the allantois of the chick embryo, purified by zonal ultracentrifugation, inactivated by formaldehyde or  $\beta$ -propiolactone, and pooled. The resulting vaccine is usually inoculated intramuscularly.

Inactivated viruses may be formulated with a suitable 15 adjuvant in order to enhance the immunological response. Such adjuvants may include but are not limited to mineral gels, e.g., aluminum hydroxide; surface active substances such as lysolecithin, pluronic polyols, polyanions; peptides; oil emulsions; and potentially useful human adjuvants such as BCG 20 and Corynebacterium parvum.

Many methods may be used to introduce the vaccine formulations described above, these include but are not limited to oral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, and intranasal routes. It may be 25 preferable to introduce the chimeric virus vaccine formulation via the natural route of infection of the pathogen for which the vaccine is designed. Where a live chimeric virus vaccine preparation is used, it may be preferable to introduce the formulation via the natural route of infection for influenza 30 virus. The ability of RSV and influenza virus to induce a vigorous secretory and cellular immune response can be used advantageously. For example, infection of the respiratory tract by chimeric RSV or influenza viruses may induce a strong secretory immune response, for example in the urogenital 35 system, with concomitant protection against a particular disease causing agent.

The following sections describe by way of example, and

not by limitation, the manipulation of the negative strand RNA viral genomes using RSV as an example to demonstrate the applicability of the methods of the present invention to generate chimeric viruses for the purposes of heterologous 5 gene expression, generating infectious viral particles and attenuated viral particles for the purposes of vaccination.

**6. RESCUE OF INFECTIOUS RESPIRATORY SYNCYTIAL VIRUSES (RSV) USING RNA DERIVED FROM SPECIFIC RECOMBINANT DNAS**

- 10 This example describes a process for the rescue of infectious respiratory syncytial virus (RSV), derived from recombinant cDNAs encoding the entire RSV RNA genome into stable and infectious RSVs, as noted in Section 5 above. The method described may be applied to both segmented and non-
- 15 segmented RNA viruses, including orthomyxovirus, paramyxovirus, e.g., Sendai virus, parainfluenza virus types 1-4, mumps, newcastle disease virus; morbillivirus, e.g., measles, canine distemper virus, rinderpest virus; pneumovirus, e.g., respiratory syncytial virus; rhabdovirus, 20 e.g., rabies, vesiculovirus, vesicular stomatitis virus; but is described by way of example in terms of RSV. This process can be used in the production of chimeric RSV viruses which can express foreign genes, i.e., genes non-native to RSV, including other viral proteins such as the HIV env protein.
- 25 Another exemplary way to achieve the production of chimeric RSV involves modifying existing, native RSV genes, as is further described. Accordingly, this example also describes the utility of this process in the directed attenuation of RSV pathogenicity, resulting in production of a vaccine with 30 defined, engineered biological properties for use in humans.

The first step of the rescue process involving the entire RSV RNA genome requires synthesis of a full length copy of the 15 kilobase (Kb) genome of RSV strain A2. This is accomplished by splicing together subgenomic double strand 35 cDNAs (using standard procedures for genetic manipulation) ranging in size from 1 kb-3.5 kb, to form the complete genomic cDNA. Determination of the nucleotide sequence of the genomic



cDNA allows identification of errors introduced during the assembly process; errors can be corrected by site directed mutagenesis, or by substitution of the error region with a piece of chemically synthesized double strand DNA. Following  
5 assembly, the genomic cDNA is positioned adjacent to a transcriptional promoter (e.g., the T7 promoter) at one end and DNA sequence which allows transcriptional termination at the other end, e.g., a specific endonuclease or a ribozyme, to allow synthesis of a plus or minus sense RNA copy of the  
10 complete virus genome *in vitro* or in cultured cells. The leader or trailer sequences may contain additional sequences as desired, such as flanking ribozyme and tandem T7 transcriptional terminators. The ribozyme can be a hepatitis delta virus ribozyme or a hammerhead ribozyme and functions to  
15 yield an exact 3' end free of non-viral nucleotides.

In accordance with this aspect of the invention, mutations, substitutions or deletions can be made to the native RSV genomic sequence which results in an increase in RSV promoter activity. Applicants have demonstrated that even  
20 an increase in RSV promoter activity greatly enhances the efficiency of rescue of RSV, allowing for the rescue of infectious RSV particles from a full-length RSV cDNA carrying the mutation. In particular, a point mutation at position 4 of the genome (C to G) results in a several fold increase in  
25 promoter activity and the rescue of infectious viral particles from a full-length RSV cDNA clone carrying the mutation.

The rescue process utilizes the interaction of full-length RSV strain A2 genome RNA, which is transcribed from the constructed cDNA, with helper RSV subgroup B virus proteins  
30 inside cultured cells. This can be accomplished in a number of ways. For example, full-length virus genomic RNA from RSV strain A2 can be transcribed *in vitro* and transfected into RSV strain B9320 infected cells, such as 293 cells using standard transfection protocols. In addition, *in vitro* transcribed  
35 genomic RNA from RSV strain A2 can be transfected into a cell line expressing the essential RSV strain A2 proteins (in the absence of helper virus) from stably integrated virus genes.

Alternatively, *in vitro* transcribed virus genome RNA (RSV strain A2) can also be transfected into cells infected with a heterologous virus (e.g., in particular vaccinia virus) expressing the essential helper RSV strain A2 proteins, specifically the N, P, L and M2/ORF1 proteins. In addition the *in vitro* transcribed genomic RNA may be transfected into cells infected with a heterologous virus, for example vaccinia virus, expressing T7 polymerase, which enables expression of helper proteins from transfected plasmid DNAs containing the helper N, P, L and M2/ORF1 genes.

As an alternative to transfection of *in vitro* transcribed genomic RNA, plasmid DNA containing the entire RSV cDNA construct may be transfected into cells infected with a heterologous virus, for example vaccinia virus, expressing the essential helper RSV strain A2 proteins and T7 polymerase, thereby enabling transcription of the entire RSV genomic RNA from the plasmid DNA containing the RSV cDNA construct. The vaccinia virus need not however, supply the helper proteins themselves but only the T7 polymerase; then helper proteins may be expressed from transfected plasmids containing the RSV N, P, L and M2/ORF1 genes, appropriately positioned adjacent to their own T7 promoters.

When replicating virus is providing the helper function during rescue experiments, the B9320 strain of RSV is used, allowing differentiation of progeny rescue directed against RSV B9320. Rescued RSV strain A2 is positively identified by the presence of specific nucleotide 'marker' sequences inserted in the cDNA copy of the RSV genome prior to rescue.

The establishment of a rescue system for native, i.e., 'wild-type' RSV strain A2 allows modifications to be introduced into the cDNA copy of the RSV genome to construct chimeric RSV containing sequences heterologous in some manner to that of native RSV, such that the resulting rescued virus may be attenuated in pathogenicity to provide a safe and efficacious human vaccine as discussed in Section 5.4 above. The genetic alterations required to cause virus attenuation may be gross (e.g., translocation of whole genes and/or

regulatory sequences within the virus genome), or minor (e.g., single or multiple nucleotide substitution(s), addition(s) and/or deletion(s) in key regulatory or functional domains within the virus genome), as further described in detail.

- 5 In addition to alteration(s) (including alteration resulting from translocation) of the RSV genetic material to provide heterologous sequence, this process permits the insertion of 'foreign' genes (i.e., genes non-native to RSV) or genetic components thereof exhibiting biological function
- 10 or antigenicity in such a way as to give expression of these genetic elements; in this way the modified, chimeric RSV can act as an expression system for other heterologous proteins or genetic elements, such as ribozymes, anti-sense RNA, specific oligoribonucleotides, with prophylactic or therapeutic
- 15 potential, or other viral proteins for vaccine purposes.

#### **6.1. RESCUE OF THE LEADER AND TRAILER SEQUENCES OF RSV STRAIN A2 USING RSV STRAIN B9320 AS HELPER VIRUS**

##### **6.1.1. VIRUSES AND CELLS**

- 20 Although RSV strain A2 and RSV strain B9320 were used in this Example, they are exemplary. It is within the skill in the art to use other strains of RSV subgroup A and RSV subgroup B viruses in accordance with the teachings of this Example. Methods which employ such other strains are
- 25 encompassed by the invention.

RSV strain A2 and RSV strain B9320 were grown in Hep-2 cells and Vero cells respectively, and 293 cells were used as host during transfection/rescue experiments. All three cell lines were obtained from the ATCC (Rockville, Maryland).

30

##### **6.1.2. CONSTRUCTION & FUNCTIONAL ANALYSIS OF REPORTER PLASMIDS**

Plasmid pRSVA2CAT (Fig. 1) was constructed as described below.

- 35 The cDNAs of the 44 nucleotide leader and 155 nucleotide trailer components of RSV strain A2 (see Mink et al., Virology

185:615-624 (1991); Collins et al., Proc. Natl. Acad. Sci. 88:9663-9667 (1991)), the trailer component also including the promoter consensus sequence of bacteriophage T7 polymerase, were separately assembled by controlled annealing of 5 oligonucleotides with partial overlapping complementarity (see Fig. 1). The oligonucleotides used in the annealing were synthesized on an Applied Biosystems DNA synthesizer (Foster City, CA). The separate oligonucleotides and their relative positions in the leader and trailer sequences are indicated in 10 Fig. 1. The oligonucleotides used to construct the leader were:

1. 5'CGA CGC ATA TTA CGC GAA AAA ATG CGT ACA ACA  
AAC TTG CAT AAA C
  2. 5'CAA AAA AAT GGG GCA AAT AAG AAT TTG ATA AGT  
ACC ACT TAA ATT TAA CT
  3. 5'CTA GAG TTA AAT TTA AGT GGT ACT
  4. 5'TAT CAA ATT CTT ATT TGC CCC ATT TTT TTG GTT  
TAT GCA AGT TTG TTG TA
  5. 5'CGC ATT TTT TCG CGT AAT ATG CGT CGG TAC
- 20 The oligonucleotides used to construct the trailer were:
1. 5'GTA TTC AAT TAT AGT TAT TAA AAA TTA AAA ATC  
ATA TAA TTT TTT AAA TA
  2. 5'ACT TTT AGT GAA CTA ATC CTA AAG TTA TCA TTT  
TAA TCT TGG AGG AAT AA
  3. 5'ATT TAA ACC CTA ATC TAA TTG GTT TAT ATG TGT  
ATT AAC TAA ATT ACG AG
  4. 5'ATA TTA GTT TTT GAC ACT TTT TTT CTC GTT ATA  
GTG AGT CGT ATT A
  5. 5'AGC TTA ATA CGA CTC ACT ATA ACG A
  6. 5'GAA AAA AAG TGT CAA AAA CTA ATA TCT CGT AAT  
TTA GTT AAT ACA CAT AT
  7. 5'AAA CCA ATT AGA TTA GGG TTT AAA TTT ATT CCT  
CCA AGA TTA AAA TGA TA
  8. 5'ACT TTA GGA TTA GTT CAC TAA AAG TTA TTT AAA  
AAA TTA TAT GAT TTT TA
  9. 5'ATT TTT AAT AAC TAT AAT TGA ATA CTG CA

The complete leader and trailer cDNAs were then ligated to the chloramphenicol-acetyl-transferase (CAT) reporter gene XbaI and PstI sites respectively to form a linear - 1 kb RSV/CAT cDNA construct. This cDNA construct was then ligated 5 into the Kpn I and Hind II sites of pUC19. The integrity of the final pRSVA2CAT construct was checked by gel analysis for the size of the Xba I/Pst I and Kpn I/Hind II digestion products. The complete leader and trailer cDNAs were also ligated to the green fluorescent protein (GFP) gene using 10 appropriate restriction enzyme sites to form a linear cDNA construct. The resulting RSV-GFP-CAT is a bicistronic reporter construct which expresses both CAT and GFP.

*In vitro* transcription of Hga I linearized pRSVA2CAT with bacteriophage T7 polymerase was performed according to 15 the T7 supplier protocol (Promega Corporation, Madison, Wisconsin). Confluent 293 cells in six-well dishes ( $1 \times 10^6$  cells per well) were infected with RSV strain B9320 at 1 plaque forming units (p.f.u.) per cell and 1 hour later were transfected with 5-10  $\mu$ g of the *in vitro* transcribed RNA from 20 the pRSVA2CAT construct. The transfection procedure followed the transfection procedure of Collins et al., Virology 195:252-256 (1993) and employed Transect/ACT™ and Opti-MEM reagents according to the manufacturers specifications (Gibco-BRL, Bethesda, Maryland). At 24 hours post-infection the 293 25 cells were assayed for CAT activity using a standard protocol (Current Protocols in Molecular Biology, Vol. 1, Chapter 9.6.2; Gorman, et al., 1982) Mol. Cell Biol. 2: 1044-1051). The detection of high levels of CAT activity indicated that *in vitro* transcribed negative sense RNA containing the 'leader' 30 and 'trailer' regions of the RSV A2 strain genome and the CAT gene can be encapsidated, replicated and expressed using proteins supplied by RSV strain B9320 (See Fig. 2). The level of CAT activity observed in these experiments was at least as high as that observed in similar rescue experiments where 35 homologous RSV strain A2 was used as helper virus. The ability of an antigenically distinct subgroup B RSV strain B9320 to support the encapsidation, replication and

transcription of a subgroup A RSV strain A2 RNA has to our knowledge hitherto not been formally reported.

## 6.2. CONSTRUCTION OF A cDNA REPRESENTING THE COMPLETE GENOME OF RSV

5 To obtain a template for cDNA synthesis, RSV genomic RNA, comprising 15,222 nucleotides, was purified from infected Hep-2 cells according to the method described by Ward et al., J. Gen. Virol. 64:167-1876 (1983). Based on the published  
10 nucleotide sequence of RSV, oligonucleotides were synthesized using an Applied Biosystems DNA synthesizer (Applied Biosystems, Foster City, CA) to act as primers for first and second strand cDNA synthesis from the genomic RNA template. The nucleotide sequences and the relative positions of the  
15 cDNA primers and key endonuclease sites within the RSV genome are indicated in Fig. 3. The production of cDNAs from virus genomic RNA was carried out according to the reverse transcription/polymerase chain reaction (RT/PCR) protocol of Perkin Elmer Corporation, Norwalk, Connecticut (see also Wang  
20 et al., (1989) Proc. Natl. Acad. Sci. 86:9717-9721); the amplified cDNAs were purified by electroelution of the appropriate DNA band from agarose gels. Purified DNA was ligated directly into the pCRII plasmid vector (Invitrogen Corp. San Diego), and transformed into either 'One Shot E.  
25 coli cells (Invitrogen) or 'SURE' E. coli cells (Stratagene, San Diego). The resulting, cloned, virus specific, cDNAs were assembled by standard cloning techniques (Sambrook et al., Molecular Cloning - A Laboratory Manual, Cold Spring Harbor laboratory Press (Cold Spring Harbor, NY, 1989) to produce a  
30 cDNA spanning the complete RSV genome. The entire cDNA genome was sequenced, and incorrect sequences were replaced by either site-directed mutagenesis or chemically synthesized DNA. Nucleotide substitutions were introduced at bases 7291 and 7294 (with base number 1 being at the start of the genomic RNA  
35 3' end) in the 'F' gene, to produce a novel Stu I endonuclease site, and at positions 7423, 7424, and 7425 (also in the F gene) to produce a novel Pme I site. These changes were

- designed to act as definitive markers for rescue events. The bacteriophage T7 polymerase and the Hga I endonuclease site were placed at opposite ends of the virus genome cDNA such that either negative or positive sense virus genome RNA can be synthesized *in vitro*. The cDNAs representing the T7 polymerase promoter sequence and the recognition sequence for Hga I were synthesized on an Applied Biosystems DNA synthesizer and were separately ligated to the ends of the virus genome cDNA, or were added as an integral part of PCR primers during amplification of the terminal portion of the genome cDNA, where appropriate; the latter procedure was used when suitable endonuclease sites near the genome cDNA termini were absent, preventing direct ligation of chemically synthesized T7 promoter/Hga I site cDNA to the genome cDNA.
- 15 This complete construct (genome cDNA and flanking T7 promoter/Hga I recognition sequence) was then cloned into the Kpn I/Not I sites of the Bluescript II SK phagemid (Stratagene, San Diego) from which the endogenous T7 promoter has been removed by site-directed mutagenesis. RNA
- 20 transcribed from this complete genome construct may be rescued using RSV subgroup B helper virus to give infectious RSV in accordance with Example 6.1. This basic rescue system for the complete native, i.e., 'wild-type' RSV A2 strain genomic RNA can be employed to introduce a variety of modifications into
- 25 the cDNA copy of the genome resulting in the introduction of heterologous sequences into the genome. Such changes can be designed to reduce viral pathogenicity without restricting virus replication to a point where rescue becomes impossible or where virus gene expression is insufficient to stimulate
- 30 adequate immunity.

The following oligonucleotides were used to construct the ribozyme/T7 terminator sequence:

5'GGT\*GGCCGGCATGGTCCCAGC  
3'CCA CCGGCCGTACCAAGGTCG

- 35 CTGCTGGCGCCGGCTGGGCAACA  
GAGCGACCCGGCCGACCCGTGTG  
TTCCGAGGGGACCGTCCCCTCGGT

AAGGCTCCCCGTGGCAGGGGAGCCA

AATGGCGAATGGGACGTCGACAGC  
TTACCGCTTACCCTGCAGCTGTGCG

TAACAAAGCCCCGAAGGAAGCT

5 ATTGTTTCGGGCTTCCTTCGA

GAGTTGCTGCTGCCACCGTTG  
CTCAACGACGACGGAGGCAAC

AGCAATAACTAGATAACCTTGGG  
TCGTTATGTGATCTATTGGAACCC

10 CCTCTAAACGGGTCTTGAGGGTCT  
GGAGATTTCGCCAGAACTCCCAGA

TTTGTCTGAAAGGAGGAACATA  
AAAACGACTTTCCTCCTTGAT

TATGCGGCCGCGTCGACGGTA

15 ATACGCCGGCGCGEAGCTGCCAT

CCGGGCCCGCCTTCGAAG3'  
GGCCCCGGCGCGAAGCTTCS'

A cDNA clone containing the complete genome of RSV a T7 promoter, a hepatitis delta virus ribozyme and a T7 terminator was generated. This construct can be used to generate antigenomic RNA or RSV *in vivo* in the presence of T7 polymerase. Sequence analysis indicated that the plasmid contained few mutations in RSV genome.

#### 25 6.2.1. MODIFICATIONS OF THE RSV GENOME

Modifications of the RSV RNA genome can comprise gross alterations of the genetic structure of RSV, such as gene shuffling. For example, the RSV M1 gene can be translocated to a position closer to the 5' end of the genome, in order to take advantage of the known 3' to 5' gradient in virus gene expression, resulting in reduced levels of M1 protein expression in infected cells and thereby reducing the rate of virus assembly and maturation. Other genes and/or regulatory regions may also be translocated appropriately, in some cases from other strains of RSV of human or animal origin. For example, the F gene (and possibly the 'G' gene) of the human



subgroup B RSV could be inserted into an otherwise RSV strain A genome (in place or, or in addition to the RSV strain A F and G genes).

- In another approach, the RNA sequence of the RSV viruses
- 5 N protein can be translocated from its 3' proximal site to a position closer to the 5' end of the genome, again taking advantage of the 3' to 5' gradient in gene transcription to reduce the level of N protein produced. By reducing the level of N protein produced, there would result a concomitant
- 10 increase in the relative rates of transcription of genes involved in stimulating host immunity to RSV and a concomitant reduction in the relative rate of genome replication. Thus, by translocating the RSV RNA sequence coding for RSV N protein, a chimeric RS virus having attenuated pathogenicity
- 15 relative to native RSV will be produced.

- Another exemplary translocation modification resulting in the production of attenuated chimeric RSV comprises the translocation of the RSV RNA sequence coding for the L protein of RSV. This sequence of the RS virus is believed responsible
- 20 for viral polymerase protein production. By translocating the RSV sequence coding for L protein from its native 5' terminal location in the native RSV genome to a location at or near the 3' terminus of the genome, a chimeric RSV virus exhibiting attenuated pathogenicity will be produced. Yet another
- 25 exemplary translocation comprises the switching the locations of the RSV RNA sequences coding for the RSV G and F proteins (i.e., relative to each other in the genome) to achieve a chimeric RSV having attenuated pathogenicity resulting from the slight modification in the amount of the G and F proteins
- 30 produced. Such gene shuffling modifications as are exemplified and discussed above are believed to result in a chimeric, modified RSV having attenuated pathogenicity in comparison to the native RSV starting material. The nucleotide sequences for the foregoing encoded proteins are
- 35 known, as is the nucleotide sequence for the entire RSV genome. See McIntosh, Respiratory Syncytial Virus in Virology, 2d Ed. edited by B.N. Fields, D.M. Knipe et al.,

al., Raven Press, Ltd. New York, 1990 Chapter 38, pp 1045-1073, and references cited therein.

- These modifications can additionally or alternatively comprise localized, site specific, single or multiple,
- 5 nucleotide substitutions, deletions or additions within genes and/or regulatory domains of the RSV genome. Such site specific, single or multiple, substitutions, deletions or additions can reduce virus pathogenicity without overly attenuating it, for example, by reducing the number of lysine
  - 10 or arginine residues at the cleavage site in the F protein to reduce efficiency of its cleavage by host cell protease (which cleavage is believed to be an essential step in functional activation of the F protein), and thereby possibly reduce virulence. Site specific modifications in the 3' or 5'
  - 15 regulatory regions of the RSV genome may also be used to increase transcription at the expense of genome replication. In addition, localized manipulation of domains within the N protein, which is believed to control the switch between transcription and replication can be made to reduce genome
  - 20 replication but still allow high levels of transcription. Further, the cytoplasmic domain(s) of the G and F glycoproteins can be altered in order to reduce their rate of migration through the endoplasmic reticulum and golgi of infected cells, thereby slowing virus maturation. In such
  - 25 cases, it may be sufficient to modify the migration of G protein only, which would then allow additional up-regulation of 'F' production, the main antigen involved in stimulating neutralizing antibody production during RSV infections. Such localized substitutions, deletions or additions within genes
  - 30 and/or regulatory domains of the RSV genome are believed to result in chimeric, modified RSV also having reduced pathogenicity relative to the native RSV genome.

### 6.3. RESCUE OF A cDNA REPRESENTING THE COMPLETE GENOME OF RSV

#### 6.3.1. THE CONSTRUCTION AND FUNCTIONAL ANALYSIS OF EXPRESSION PLASMIDS

5 The RSV, N, P, and L genes encode the viral polymerase of RSV. The function of the RSV M genes is unknown. The ability of RSV, N, P, M, and L expression plasmids to serve the function of helper RSV strain AZ proteins was assessed as described below. The RSV, N, P, L, and M2-1 genes were cloned  
10 into the modified PCITE 2a(+) vector (Novagen, Madison, WI) under the control of the T7 promoter and flanked by a T7 terminator at it's 3' end. PCITE-2a(+) was modified by insertion of a T7 terminator sequence from PCITE-3a(+) into the Alwn I and Bgl II sites of pCITE-2a(+). The functionality  
15 of the N, P, and L expression plasmids was determined by their ability to replicate the transfected pRSVA2CAT. At approximately 80% confluency, Hep-2 cells in six-well plates were infected with MVA at a moi of 5. After 1 hour, the infected cells were transfected with pRSVA2CAT (0.5mg), and  
20 plasmids encoding the N (0.4mg), P (0.4mg), and L (0.2mg) genes using lipofectACE (Life Technologies, Gaithersburg, M.D.). The transfection proceeded for 5 hours or overnight and then the transfection medium was replaced with fresh MEM containing 2% (fetal bovine serum) FBS. Two days post-  
25 infection, the cells were lysed and the lysates were analyzed for CAT activity using Boehringer Mannheim's CAT ELISA kit. CAT activity was detected in cells that had been transfected with N, P, and L plasmids together with pRSVAZCAT. However, no CAT activity was detected when any one of the expression  
30 plasmids was omitted. Furthermore, co-transfection of RSV-GFP-CAT with the N, P, and L expression plasmids resulted in expression of both GFP and CAT proteins. The ratios of different expression plasmids and moi of the recombinant vaccina virus were optimized in the reporter gene expression  
35 system.

### 6.3.2. RECOVERY OF INFECTIOUS RSV FROM THE COMPLETE RSV cDNA

Hep-2 cells were infected with MVA (recombinant vaccinia virus expressing T7 polymerase) at an moi of one. Fifty minutes later, transfection mixture was added onto the cells. The transfection mixture consisted of 2  $\mu$ g of N expression vector, 2  $\mu$ g of P expression vector, 1  $\mu$ g of L expression vector, 1.25  $\mu$ g of M2/orf1 expression vector, 2  $\mu$ g of RSV genome clone with enhanced promoter, 50  $\mu$ l of LipofectACE (Life Technologies, Gaithersburg, M.D.) and 1 ml OPTI-MEM. One day later, the transfection mixture was replaced by MEM containing 2% FCS. The cells were incubated at 37°C for 2 days. The transfection supernatant was harvested and used to infect fresh Hep-2 cells in the presence of 40  $\mu$ g/ml arac (drug against vaccinia virus). The infected Hep2 cells were incubated for 7 days. After harvesting the P1 supernatant, cells were used for immunostaining using antibodies directed against F protein of RSV A2 strain. Six positively stained loci with visible cell-cell-fusion (typical for RSV infection) were identified. The RNA was extracted from P1 supernatant, and used as template for RT-PCR analysis. PCR products corresponding to F and M2 regions were generated. both products contained the introduced markers. In control, PCR products derived from natural RSV virus lacked the markers.

A point mutation was created at position 4 of the leader sequence of the RSV genome clone (C residue to G) and this genome clone was designated pRSVC4GLwt. This clone has been shown in a reporter gene context to increase the promoter activity by several fold compared to wild-type. After introduction of this mutation into the full-length genome, infectious virus was rescued from the cDNA clone. The rescued recombinant RSV virus formed smaller plaques than the wild-type RSV virus (Figure 8).

This system allows the rescue mutated RSV. Therefore, it may be an excellent tool to engineer live-attenuated vaccines against RSV and to use RSV vector and viruses to achieve heterologous gene expression. It may be possible to express G

protein of type B RSV into the type A background, so the vaccine is capable of protect both type A and type B RSV infection. It may also be possible to achieve attenuation and temperature sensitive mutations into the RSV genome, by 5 changing the gene order or by site-directed mutagenesis of the L protein.

#### 6.4. USE OF MONOCLONAL ANTIBODIES TO DIFFERENTIATE RESCUED VIRUS FROM HELPER VIRUS

10 In order to neutralize the RSV strain B9320 helper virus and facilitate identification of rescued A 2 strain RSV, monoclonal antibodies against RSV strain B9320 were made as follows.

Six BALB/c female mice were infected intranasally (i.n.) with  $10^5$  plaque forming units (p.f.u.) of RSV B9320, followed 15 5 weeks later by intraperitoneal (i.p.) inoculation with  $10^6$ - $10^7$  pfu of RSV B9320 in a mixture containing 50% complete Freund's adjuvant. Two weeks after i.p. inoculation, a blood sample from each mouse was tested for the presence of RSV specific antibody using a standard neutralization assay 20 (Beeler and Coelingh, J. Virol. 63:2941-2950 (1988)). Mice producing the highest level of neutralizing antibody were then further boosted with  $10^6$  p.f.u. of RSV strain B9320 in phosphate buffered saline (PBS), injected intravenously at the 25 base of the tail. Three days later, the mice were sacrificed and their spleens collected as a source of monoclonal antibody producing B-cells. Splenocytes (including B-cells) were teased from the mouse spleen through incisions made in the spleen capsule into 5 ml of Dulbecco's Modified Eagle's Medium (DME). Clumps of cells were allowed to settle out, and the 30 remaining suspended cells were separately collected by centrifugation at 2000xg for 5 minutes at room temperature. These cell pellets were resuspended in 15 ml 0.83 (W/V)  $\text{NH}_4\text{Cl}$ , and allowed to stand for 5 minutes to lyse red blood cells. 35 Splenocytes were then collected by centrifugation as before through a 10 ml; cushion of fetal calf serum. The splenocytes were then rinsed in DME, repelleted and finally resuspended in

20 ml of fresh DME. These splenocytes were then mixed with Sp2/0 cells (a mouse myeloma cell line used as fusion partners for the immortalization of splenocytes) in a ratio of 10:1, spleen cells: Sp2/0 cells. Sp2/0 cells were obtained from the 5 ATCC and maintained in DME supplemented with 10% fetal bovine serum. The cell mixture was then centrifuged for 8 minutes at 2000xg at room temperature. The cell pellet was resuspended in 1 ml of 50% polyethylene glycol 1000 mol. wt. (PEG 1000), followed by addition of equal volumes of DME at 1 minute 10 intervals until a final volume of 25 ml was attained. The fused cells were then pelleted as before and resuspended at  $3.5 \times 10^6$  spleen cells  $\text{ml}^{-1}$  in growth medium (50% conditioned medium from SP2/0 cells, 50% HA medium containing 100 ml RPMI 25 ml F.C.S., 100  $\mu\text{g/ml}$  gentamicin, 4 ml 50X Hypoxanthine, 15 Thymidine, Aminopterin (HAT) medium supplied as a prepared mixture of Sigma Chem. Co., St. Louis, MO). The cell suspension was distributed over well plates (200  $\mu\text{l}$  well $^{-1}$ ) and incubated at 37°C, 95 humidity and 5% CO<sub>2</sub>. Colonies of hybridoma cells (fused splenocytes and Sp2/0 cells) were then 20 subcultured into 24 well plates and grown until nearly confluent; the supernatant growth medium was then sampled for the presence of RSV strain B9320 neutralizing monoclonal antibody, using a standard neutralization assay (Beeler and Coelingh, J. Virol. 63:2941-50 (1988)). Hybridoma cells from 25 wells with neutralizing activity were resuspended in growth medium and diluted to give a cell density of 0.5 cells per 100  $\mu\text{l}$  and plated out in 96 well plates, 200  $\mu\text{l}$  per well. This procedure ensured the production of monoclones (i.e. hybridoma cell lines derived from a single cell) which were then 30 reassayed for the production of neutralizing monoclonal antibody. Those hybridoma cell lines which produced monoclonal antibody capable of neutralizing RSV strain B9320 but not RSV strain A2 were subsequently infected into mice, i.p. ( $10^6$  cells per mouse). Two weeks after the i.p. 35 injection mouse ascites fluid containing neutralizing monoclonal antibody for RSV strain B9320 was tapped with a 19 gauge needle, and stored at -20°C.

This monoclonal antibody was used to neutralize the RSV strain B9320 helper virus following rescue of RSV strain A2 as described in Section 9.1. This was carried out by diluting neutralizing monoclonal antibody 1 in 50 with molten 0.4% (w/v) agar in Eagle's Minimal Essential Medium (EMEM) containing 1% F.C.S. This mixture was then added to Hep-2 cell monolayers, which had been infected with the progeny of rescue experiments at an m.o.i. of 0.1-0.01 p.f.u. per cell. The monoclonal antibody in the agar overlay inhibited the growth of RSV strain B9320, but allowed the growth of RSV strain A2, resulting in plaque formation by the A2 strain. These plaques were picked using a pasteur pipette to remove a plug of agar above the plaque and the infected cells within the plaque; the cells and agar plug were resuspended in 2 ml of EMEM, 1% FCS, and released virus was plaqued again in the presence of monoclonal antibody on a fresh Hep-2 cell monolayer to further purify from helper virus. The twice plaqued virus was then used to infect Hep-2 cells in 24 well plates, and the progeny from that were used to infect six-well plates at an m.o.i. of 0.1 p.f.u. per cell. Finally, total infected cell RNA from one well of a six-well plates was used in a RT/PCR reaction using first and second strand primers on either side of the 'marker sequences' (introduced into the RSV strain A2 genome to act as a means of recognizing rescue events) as described in Section 6.2 above. The DNA produced from the RT/PCR reaction was subsequently digested with Stu I and Pme I to positively identify the 'marker sequences' introduced into RSV strain A2 cDNA, and hence to establish the validity of the rescue process.

30

#### 7. RESCUE OF INFECTIOUS RSV PARTICLES IN THE ABSENCE OF M2 EXPRESSION

The following experiments were conducted to compare the efficiencies of rescue of RS virions in the presence and absence of the M2/ORF1 gene. If the M2/ORF1 gene function is not required to achieve rescue of RSV infectious particles, it should be possible to rescue RS virions in the absence of the

35

expression of the M2/ORF1 gene function. In the present analysis, Hep-2 cells which are susceptible to RSV replication, were co-transfected with plasmids encoding the 'N', 'P' and 'L' genes of the viral polymerase of RSV and the 5 cDNA corresponding to the full-length antigenome of RSV, in the presence or absence of plasmid DNA encoding the M2/ORF1 gene, and the number of RSV infectious units were measured in order to determine whether or not the M2/ORF1 gene product was required to rescue infectious RSV particles.

- 10 The following plasmids were used in the experiments described below: a cDNA clone encoding the full-length antigenome of RSV strain A2, designated pRSVC4GLwt; and plasmids encoding the N, P, and L polymerase proteins, and plasmid encoding the M2/ORF1 elongation factor, each 15 downstream of a T7 RNA promoter, designated by the name of the viral protein encoded.

- pRSVC4GLwt was transfected, together with plasmids encoding proteins N, P and L, into Hep-2 cells which had been pre-infected with a recombinant vaccinia virus expressing the 20 T7 RNA polymerase (designated MVA). In another set of Hep-2 cells, pRSVC4GLwt was co-transfected with plasmids encoding the N, P and L polymerase proteins, and in addition a plasmid encoding the M2 function. Transfection and recovery of recombinant RSV were performed as follows: Hep-2 cells were 25 split in six-well dishes (35mm per well) 5 hours or 24 hours prior to transfection. Each well contained approximately  $1 \times 10^6$  cells which were grown in MEM (minimum essential medium) containing 10% FBS (fetal bovine serum). Monolayers of Hep-2 cells at 70% - 80% confluence were infected with MVA at a 30 multiplicity of infection (moi) of 5 and incubated at 35°C for 60 minutes. The cells were then washed once with OPTI-MEM (Life Technologies) and the medium of each dish replaced with 1 ml of OPTI-MEM and 0.2 ml of the transfection mixture. The transfection mixture was prepared by mixing the four plasmids, 35 pRSVC4GLwt, N, P and L plasmids in a final volume of 0.1 ml OPTI-MEM at amounts of 0.5 - 0.6  $\mu$ g of pRSVC4GLwt, 0.4  $\mu$ g of N plasmid, 0.4  $\mu$ g of P plasmid, and 0.2  $\mu$ g of L plasmid. A



second mixture was prepared which additionally included 0.4  $\mu$ g M2/ORFI plasmid. The plasmid mixtures of 0.1 ml were combined with 0.1 ml of OPTI-MEM containing 10  $\mu$ l of lipofectACE (Life Technologies, Gaithersburg, M.D.) to constitute the complete 5 transfection mixture. After a 15 minute incubation at room temperature, the transfection mixture was added to the cells, and one day later this was replaced by MEM containing 2% FBS. Cultures were incubated at 35°C for 3 days at which time the supernatants were harvested. Cells were incubated at 35°C 10 since the MVA virus is slightly temperature sensitive and is much more efficient at 35°C.

Three days post-transfection, the transfected cell supernatants were assayed for the presence of RSV infectious units by an immunoassay which would indicate the presence of 15 RSV packaged particles (see Table I). In this assay, 0.3 - 0.4 ml of the culture supernatants were passaged onto fresh (uninfected) Hep-2 cells and overlaid with 1% methylcellulose and 1 x L15 medium containing 2% FBS. After incubation for 6 days, the supernatant was harvested and the cells were fixed 20 and stained by an indirect horseradish peroxidase method, using a goat anti-RSV antibody which recognizes the RSV viral particle (Biogenesis, Sandown, NH) followed by a rabbit anti-goat antibody conjugated to horseradish peroxidase. The antibody complexes that bound to RSV-infected cells were 25 detected by the addition of a AEC-(3-amino-9-ethylcarbazole) chromogen substrate (DAKO) according to the manufacturer's instructions. The RSV plaques were indicated by a black-brown coloration resulting from the reaction between the chromogen substrate and the RSV-antibody complexes bound to the plaques. 30 The number of RSV plaques is expressed as the number of plaque forming units (p.f.u.) per 0.5 ml of transfection supernatant (see Table I).

Comparisons of the amount of RS virions recovered from the supernatants of transfection dishes in the presence or 35 absence of M2/ORFI are shown in Table I. The results of four separate experiments demonstrated that the absence of M2/ORFI from the transfection assay did not diminish the number of

infectious units of RSV observed. Thus, the results of these experiments clearly indicate that RSV can be rescued in the absence of the M2/ORF1 from cells transfected only with plasmids encoding the three polymerase proteins, N, P and L, 5 and the cDNA encoding the full-length RSV antigenome. The rescue of true RS virions in the absence of M2/ORF1 was further indicated by the ability to passage the rescued recombinant RSV for up to six passages. Therefore, the production of RSV virions is not dependent on the expression 10 of the M2/ORF1 gene, nor does the inclusion of the M2/ORF1 gene in the transfection assay increase the efficiency of true RSV rescue.

Table I. Production of infectious RSV through plasmid transfection is not dependent on expression of M2ORF1

15	Expt.	Production of infectious RSV (pfu from 0.5ml transfection supernatants)	
		+ M2 ORF1	-M2 ORF1
	1.	6,10(8)	16,9(13)
	2.	120,46,428(198)	100,122,105(109)
20	3.	160,180(170)	150,133(142)
	4.	588,253,725(522)	300,1000,110(470)

Each experiment was done singly, in duplicates or triplicates. The average number of plaque forming units (pfu) from 0.5 ml transfected cell supernatants is shown in the brackets.

## 25 8. EXAMPLE: EXPRESSION OF RSV SUBGROUP B-G AND -F PROTEINS BY RSV A2 STRAIN

The following experiments were conducted to generate a chimeric RSV which expresses the antigenic polypeptides of more than one strain of RSV. Two main antigenic subgroups (A and B) of respiratory syncytial virus (RSV) cause human 30 diseases. Glycoproteins F and G are the two major antigenic determinants of RSV. The F glycoproteins of subgroup A and B viruses are estimated to be 50% related, while the relationship of G glycoproteins is considerably less, about 1- 35 5%. Infection of RSV subgroup A induces either partial or no resistance to replication of a subgroup B strain and vice

versa. Both subgroup A and subgroup B RSV virus vaccines are needed to protect from RSV infection.

The first approach described herein is to make an infectious chimeric RSV cDNA clone expressing subgroup B antigens by replacing the current infectious RSV A2 cDNA clone G and F region with subgroup B-G and -F genes. The chimeric RSV would be subgroup B antigenic specific. The second approach described herein is to insert subgroup B-G gene in the current A2 cDNA clone so that one virus would express both subgroup A and B specific antigens.

### 8.1. Substitution of A2 G and F by B9320 G and F genes

RSV subgroup B strain B9320 G and F genes were amplified from B9320 vRNA by RT/PCR and cloned into pCRII vector for sequence determination. BamH I site was created in the oligonucleotide primers used for RT/PCR in order to clone the G and F genes from B9320 strain into A2 antigenomic cDNA (Fig. 4A). A cDNA fragment which contained G and F genes from 4326 nt to 9387 nt of A2 strain was first subcloned into pUC19 (pUCR/H). Bgl II sites were created at positions of 4630 (SH/G intergenic junction) and 7554 (F/M2 intergenic junction), respectively by Quickchange site-directed mutagenesis kit (Stratagene, La Jolla, CA). B9320 G and F cDNA inserted in pCR.II vector was digested with BamH I restriction enzyme and then subcloned into Bgl II digested pUCR/H which had the A2 G and F genes removed. The cDNA clone with A2 G and F genes replaced by B9320 G and F was used to replace the Xho I to Msc I region of the full-length A2 antigenomic cDNA. The resulting antigenomic cDNA clone was termed pRSVB-GF and was used to transfect Hep-2 cells to generate infectious RSVB-GF virus.

Generation of chimeric RSVB-GF virus was as follows, pRSVB-GF was transfected, together with plasmids encoding proteins N, P, L and M2/ORF1, into Hep-2 cells which had been infected with MVA, a recombinant vaccinia virus which expresses the T7 RNA polymerase. Hep-2 cells were split a day before transfection in six-well dishes. Monolayers of Hep-2

cells at 60% - 70% confluence were infected with MVA at moi of 5 and incubated at 35°C for 60 min. The cells were then washed once with OPTI-MEM (Life Technologies, Gaithersburg, MD). Each dish was replaced with 1 ml of OPTI-MEM and added 5 with 0.2 ml of transfection medium. The transfection medium was prepared by mixing five plasmids in a final volume of 0.1 ml of OPTI-MEM medium, namely 0.6 µg of RSV antigenome pRSVB-GF, 0.4 µg of N plasmid, 0.4 µg of P plasmid, 0.2 µg of L plasmid and 0.4 µg of M2/ORF1 plasmid. This was combined 10 with 0.1 ml of OPTI-MEM containing 10 µl lipofectACE (Life Technologies, Gaithersburg, MD U.S.A.). After a 15 minute incubation at room temperature, the DNA/lipofectACE was added to the cells and the medium was replaced one day later by MEM containing 2% FBS. Cultures were further incubated at 35°C 15 for 3 days and the supernatants harvested. Aliquots of culture supernatants (PO) were then used to infect fresh Hep-2 cells. After incubation for 6 days at 35°C, the supernatant was harvested and the cells were fixed and stained by an indirect horseradish peroxidase method using goat anti-RSV 20 antibody (Biogenesis, Sandown, NH) followed by a rabbit anti-goat antibody linked to horseradish peroxidase. The virus infected cells were then detected by addition of substrate chromogen (DAKO, Carpinteria, CA, U.S.A.) according to the manufacturer's instructions. RSV-like plaques were detected 25 in the cells which were infected with the supernatants from cells transfected with pRSVB-GF. The virus was further plaque purified twice and amplified in Hep-2 cells.

Recombinant RSVB-GF virus was characterized by RT/PCR using RSV subgroup B specific primers. Two independently 30 purified recombinant RSVB-GF virus isolates were extracted with an RNA extraction kit (Tel-Test, Friendswood, TX) and RNA was precipitated by isopropanol. Virion RNAs were annealed with a primer spanning the RSV region from nt 4468 to 4492 and 35 reactions) using superscript reverse transcriptase (Life Technologies, Gaithersburg, MD). Aliquots of each reaction were subjected to PCR (30 cycles at 94°C for 30 s, 55°C for

30 s and 72°C for 2 min) using subgroup B specific primers in G region (CACCACCTACTTTACTCAAGT and TTGTTTGTGGGTTTGATGGTTGG). The PCR products were analyzed by electrophoresis on 1% agarose gel and visualized by staining with ethidium bromide.

5 As shown in Fig. 5, no DNA product was produced in RT/PCR reactions using RSV A2 strain as template. However, a predicted product of 254 bp was detected in RT/PCR reactions utilizing RSVB-GF RNA or the PCR control plasmid, pRSVB-GF DNA, as template, indicating the rescued virus contained G and

10 F genes derived from B9320 virus.

### 8.2. Expression of B9320G by RSV A2 virus

RSV subgroup B strain B9320 G gene was amplified from B9320 vRNA by RT/PCR and cloned into pCRII vector for sequence

15 determination. Two Bgl II sites were incorporated into the PCR primers which also contained gene start and gene end signals (GATATCAAGATCTACAATAACATTGGGGCAAATGC and GCTAAGAGATCTTTTGAATAACTAAGCATG). B9320G cDNA insert was digested with Bgl II and cloned into the SH/G (4630 nt) or

20 F/M2 (7552 nt) intergenic junction of a A2 cDNA subclone (Fig. 4B and Fig. 4C). The Xho I to Msc I fragment containing B9320G insertion either at SH/G or F/M2 intergenic region was used to replace the corresponding Xho I to Msc I region of the A2 antigenomic cDNA. The resulting RSV antigenomic cDNA clone

25 was termed as pRSVB9320G-SH/G or pRSVB9320G-F/M2.

Generation of RSV A2 virus which had B9320 G gene inserted at F/M2 intergenic region was performed similar to what has described for generation of RSVB-GF virus. Briefly, pRSVB9320G-F/M2 together with plasmids encoding proteins N, P

30 and L were transfected, into Hep-2 cells, infected with a MVA vaccinia virus recombinant, which expresses the T7 RNA polymerase (Life Technologies, Gaithersburg, M.D.). The transfected cell medium was replaced by MEM containing 2% fetal bovine serum (FBS) one day after transfection and

35 further incubated for 3 days at 35°C. Aliquots of culture supernatants (PO) were then used to infect fresh Hep-2 cells. After incubation for 6 days at 35°C, the supernatant was

harvested and the cells were fixed and stained by an indirect horseradish peroxidase method using goat anti-RSV antibody (Biogenesis) followed by a rabbit anti-goat antibody linked to horseradish peroxidase. The virus infected cells were then  
5 detected by addition of substrate chromogen (Dako). RSV-like plaques were detected in the cells which were infected with the supernatants from cells transfected with pRSVB9320G/F/M2.

Characterization of pRSVB9320G-F/M2 virus was performed by RT/PCR using B9320G specific primers. A predicted PCR  
10 product of 410bp was seen in RT/PCR sample using pRSVB9320G-F/M2 RNA as template, indicating the rescued virus contained G gene derived from B9320. (Figure 6)

Expression of the inserted RSV B9320 G gene was analyzed by Northern blot using a <sup>32</sup>P-labeled oligonucleotide specific  
15 to A2-G or B-G mRNA. Total cellular RNA was extracted from Hep-2 cells infected with wild-type RSVB 9320, rRSVA2, or rRSVB9320G-F/M2 48 hours postinfection using an RNA extraction kit (RNA stat-60, Tel-Test). RNA was electrophoresed on a 1.2% agarose gel containing formaldehyde and transferred to a  
20 nylon membrane (Amersham). An oligonucleotide specific to the G gene of the A2 strain (5'TCTTGACTGTTGTGGATTGCAGGGTTGACTTGACTCCGATCGATCC-3') and an oligonucleotide specific to the B9320 G gene (5'CTGTGTGTTGTTGTGTATGGTGTGTTTCTGATTTTGTATTGATCGATCC-3') were  
25 labeled with <sup>32</sup>P-ATP by a kinasing reaction known to those of ordinary skill in the art. Hybridization of the membrane with one of the <sup>32</sup>P-labeled G gene specific oligonucleotides was performed at 65°C and washed according to standard procedure. Both A2-G and B9320-G specific RNA were detected in the  
30 rRSVB9320G-F/M2 infected Hep-2 Cells. (Figure 6B) These results demonstrate subtype specific RNA expression.

Protein expression of the chimeric rRSVA2(B-G) was compared to that of RSV B9320 and rRSV by immunoprecipitation of <sup>35</sup>S-labeled infected Hep-2 cell lysates. Briefly, the  
35 virus infected cells were labeled with <sup>35</sup>S-promix (100 µCi/ml <sup>35</sup>S-Cys and <sup>35</sup>S-Met, Amersham, Arlington Heights, IL) at 14 hours to 18 hours post-infection according to a protocol known

to those of ordinary skill in the art. The cell monolayers were lysed by RIPA buffer and the polypeptides were immunoprecipitated with either polyclonal antiserum raised in goat against detergent disrupted RSV A2 virus (Fig. 7, lanes 1-4) or antiserum raised in mice against undisrupted B9320 virions (Fig. 7, lanes 5-8). The radio labeled immunoprecipitated polypeptides were electrophoresed on 10% polyacrylamide gels containing 0.1% SDS and detected by autoradiography. Anti-RSV A2 serum immunoprecipitated the major polypeptides of the RSV A2 strain, whereas anti-B9320 serum mainly reacted with RSV B9320 G protein and the conserved F protein of both A and B subgroups. As shown in Fig. 7, a protein which is identical to the A2-G protein (lane 3), was immunoprecipitated from the rRSVA2(B-G) infected cells (lane 4) by using an antiserum against RSV A2. The G protein of RSV B9320 strain was not recognized by the anti-A2 antiserum. A protein species, smaller than A2-G protein, was immunoprecipitated from both B9320 (lane 6) and rRSVA2(B-G) (lane 9) infected cells using the antiserum raised in mice against B9320 virions. This polypeptide was not present in the uninfected and RSV A2 infected cells and likely is to represent the G protein specific to the RSV B 9320 strain. Amino acid sequence comparison of both A2 and B9320 RSV G proteins indicated that two additional potential N-glycosylation sites (N-X-S/t) are present in the RSV A2G protein, which may contribute to slower migration of the A2 G protein under the conditions used. The F protein of RSV B9320 also migrated slightly faster than RSV A2 F protein. The P and M proteins also showed mobility differences between the two virus subtypes. The identity of the polypeptide near the top of the protein gel present in FSV B9320 and rRSVA2(B-G) infected cells is not known. Antisera raised in mice of RSV B9320 virions poorly recognized the N, P and M proteins are compared to the goat antiserum raised against the RSV A2 strain. The data described above clearly indicate that chimeric rRSV A2(B-G) expresses both the RSV A2 and B9320 specific G proteins.

### 8.2.1 Replication of Recombinant RSV in Tissue Culture

Recombinant RS viruses were plaque purified three times and amplified in Hep-2 cells. Plaque assays were performed in Hep-2 cells in 12-well plates using an overlay of 1%

- 5 methylcellulose and 1 x L15 medium containing 2% fetal bovine serum (FBS). After incubation at 35°C for 6 days, the monolayers were fixed with methanol and plaques were identified by immunostaining. Plaque size and morphology of rRSV was very similar to that of wild-type A2 RSV (Fig. 8).
- 10 However, the plaques formed by rRSVC4G were smaller than rRSV and wild-type A2 virus. The only genetic difference between rRSV and rRSVC4 was a single nucleotide substitution in the RSV leader region. Therefore, the smaller plaque size of rRSV A2(B-G) was not distinguishable from that of rRSVC4G.
- 15 The growth curves of rRSV, rRSVC4G and rRSV A2 (B-G) were compared to that of the biologically derived wild-type A2 virus. Hep-2 cells were grown in T25 culture flasks and infected with rRSV, rRSVC4G, rRSVA2(B-G), or wild-type RSV A2 strain at a moi of 0.5. After 1 hour adsorption at 37°C, the
- 20 cells were washed three times with MEM containing 2% FBS and incubated at 37°C in 5% CO<sub>2</sub>. At 4 hour intervals post-infection, 250 µl of the culture supernatant was collected, and stored at -70°C until virus titration. Each aliquot taken was replaced with an equal amount of fresh medium. The titer
- 25 of each virus was determined by plaque assay on Hep-2 cells and visualized by immunostaining (*vide supra*). As shown in Fig. 9, the growth kinetics of rRSV is very similar to that of wild-type A2 virus. Maximum virus titer for all the viruses were achieved between 48 hr to 72 hr. The virus titer of
- 30 rRSVC4G was about 2.4-fold (at 48 hr) and 6.6-fold (at 72 hr) lower than rRSV and wild-type A2 RSV. The poor growth of rRSVC4G may also be due to the single nucleotide change in the leader region. The chimeric rRSV A2(B-G) showed slower kinetics and lower peak titer (Fig. 9).

35



## 9. EXAMPLE: GENERATION OF RSV L GENE MUTANTS

The strategy for generating L gene mutants is to introduce defined mutations or random mutations into the RSV L gene. The functionality of the L gene cDNA mutants can be  
5 screened *in vitro* by a minigenome replication system. The recovered L gene mutants are then further analyzed *in vitro* and *in vivo*.

### 9.1 MUTAGENESIS STRATEGIES

10

#### 9.1.1 SCANNING MUTAGENESIS TO CHANGE THE CLUSTERED CHARGED AMINO ACIDS TO ALANINE

This mutagenesis strategy has been shown to be particularly effective in systematically targeting functional  
15 domains exposed on protein surfaces. The rationale is that clusters of charged residues generally do not lie buried in the protein structure. Making conservative substitutions of these charged residues with alanines will therefore remove the charges without grossly changing the structure of the protein.  
20 Disruption of charged clusters may interfere with the interaction of RSV L protein with other proteins and make its activity thermosensitive, thereby yielding temperature-sensitive mutants.

A cluster was originally defined arbitrarily as a stretch  
25 of 5 amino acids in which two or more residues are charged residues. For scanning mutagenesis, all the charged residues in the clusters can be changed to alanines by site directed mutagenesis. Because of the large size of the RSV L gene, there are many clustered charged residues in the L protein.  
30 Therefore, only contiguous charged residues of 3 to 5 amino acids throughout the entire L gene were targeted (Fig. 10). The RSV L protein contains 2 clusters of five contiguous charged residues, 2 clusters of four contiguous charged residues and 17 clusters of three contiguous charge residues.  
35 Two to four of the charged residues in each cluster were substituted with alanines.

The first step of the invention was to introduce the changes into pCITE-L which contains the entire RSV L-gene, using a QuikChange site-directed mutagenesis kit (Stratagene). The introduced mutations were then confirmed by sequence analysis.

#### 9.1.2. CYSTEINE SCANNING MUTAGENESIS

Cysteines are good targets for mutagenesis as they are frequently involved in intramolecular and intermolecular bond formations. By changing cysteines to glycines or alanines, the stability and function of a protein may be altered because of disruption of its tertiary structure. Thirty-nine cysteine residues are present in the RSV L protein (Fig. 11). Comparison of the RSV L protein with other members of paramyxoviruses indicates that some of the cysteine residues are conserved.

Five conserved cysteine residues were changed to either valine (conservative change) or to aspartic acids (nonconservative change) using a QuikChange site-directed mutagenesis kit (Stratagene) degenerate mutagenic oligonucleotides. It will be apparent to one skilled in the art that the sequence of the mutagenic oligonucleotides is determined by the protein sequence desired. The introduced mutations were confirmed by sequence analysis.

25

#### 9.1.3. RANDOM MUTAGENESIS

Random mutagenesis may change any residue, not simply charged residues or cysteines. Because of the size of the RSV L gene, several L gene cDNA fragments were mutagenized by PCR mutagenesis. This was accomplished by PCR using exo Pfu polymerase obtained from Stratagene. Mutagenized PCR fragments were then cloned into a pCITE-L vector. Sequencing analysis of 20 mutagenized cDNA fragments indicated that 80% - 90% mutation rates were achieved. The functionality of these mutants was then screened by a minigenome replication system. Any mutants showing altered polymerase function were then

further cloned into the full-length RSV cDNA clone and virus recovered from transfected cells.

### 9.2. FUNCTIONAL ANALYSIS OF RSV L PROTEIN MUTANTS BY MINIGENOME REPLICATION SYSTEM

5 The functionality of the L-genes mutants were tested by their ability to replicate a RSV minigenome containing a CAT gene in its antisense and flanked by RSV leader and trailer sequences. Hep-2 cells were infected with MVA vaccinia  
10 recombinants expressing T7 RNA polymerase. After one hour, the cells were transfected with plasmids expressing mutated L protein together with plasmids expressing N protein and P protein, and pRSV/CAT plasmid containing CAT gene (minigenome). CAT gene expression from the transfected cells  
15 was determined by a CAT ELISA assay (Boehringer Mannheim) according to the manufacturer's instruction. The amount of CAT activity produced by the L gene mutant was then compared to that of wild-type L protein.

### 9.3. RECOVERY OF MUTANT RECOMBINANT RSV

20 To recover or rescue mutant recombinant RSV, mutations in the L-gene were engineered into plasmids encoding the entire RSV genome in the positive sense (antigenome). The L gene cDNA restriction fragments (BamH I and Not I) containing  
25 mutations in the L-gene were removed from pCITE vector and cloned into the full-length RSV cDNA clone. The cDNA clones were sequenced to confirm that each contained the introduced mutations.

Each RSV L gene mutant virus was rescued by co-  
30 transfection of the following plasmids into subconfluent Hep-2 cells grown in six-well plates. Prior to transfection, the Hep-2 cells were infected with MVA, a recombinant vaccinia virus which expresses T7 RNA polymerase. One hour later, cells were transfected with the following plasmids:

- 35
- pCITE-N: encoding wild-type RSV N gene, 0.4  $\mu$ g
  - pCITE-P: encoding wild-type RSV P gene, 0.4  $\mu$ g

- pCITE-Lmutant: encoding mutant RSV L gene, 0.2  $\mu$ g
- pRSVL mutant: full-length genomic RSV of the positive sense (antigenome) containing the same L-gene mutations as pCITE-L mutant, 0.6  $\mu$ g

DNA was introduced into cells by lipofectACE (Life Technologies) in OPTI-MEM. After five hours or overnight transfection, the transfection medium was removed and replaced with 2% MEM. Following incubation at 35°C for three days, the media supernatants from the transfected cells were used to infect Vero cells. The virus was recovered from the infected Vero cells and the introduced mutations in the recovered recombinant viruses confirmed by sequencing of the RT/PCR DNA derived from viral RNA.

Examples of the L gene mutants obtained by charged to alanine scanning mutagenesis are shown in the Table II. Mutants were assayed by determining the expression of CAT by pRSV/CAT minigenome following co-transfection of plasmids expressing N, P and either wild-type or mutant L. Cells were harvested and lysed 40 hours post-transfection after incubation at 33°C or 39°C. The CAT activity was monitored by CAT ELISA assay (Boehringer Mannheim). Each sample represents the average of duplicate transfections. The amount of CAT produced for each sample was determined from a linear standard curve.

From the above preliminary studies, different types of mutations have been found.

#### 9.3.1. DETRIMENTAL MUTATIONS

Seven L protein mutants displayed a greater than 99% reduction in the amount of CAT produced compared to that of wild-type L protein. These mutations drastically reduced the activity of the RSV polymerase and are not expected to be viable.

### 9.3.2. INTERMEDIATE MUTATIONS

Several L mutants showed an intermediate level of CAT production which ranged from 1% to 50% of that wild-type L protein. A subset of these mutants were introduced into virus  
5 and found to be viable. Preliminary data indicated that mutant A2 showed 10-to 20-fold reduction in virus titer when grown at 40°C compared 33°C. Mutant A25 exhibited a smaller plaque formation phenotype when grown at both 33°C and 39°C. This mutant also had a 10-fold reduction in virus titer at  
10 40°C compared to 33°C.

### 9.3.3. MUTANTS WITH L PROTEIN FUNCTION SIMILAR OR HIGHER THAN WILD TYPE L PROTEIN

Some L gene mutants produced CAT gene expression levels  
15 similar to or greater than the wild-type L protein in vitro and the recovered virus mutants have phenotypes indistinguishable from wild-type viruses in tissue culture.

Once mutations in L that confer temperature sensitivity and attenuation have been identified, the mutations will be  
20 combined to rest for the cumulative effect of multiple temperature-sensitivity markers. The L mutants bearing more than one temperature sensitive marker are expected to have lower permissive temperature and to be genetically more stable than single-marker mutants.

The generated L gene mutants may also be combined with  
25 mutations present in other RSV genes and/or with non-essential RSV gene deletion mutants (e.g., SH and M2-2 deletion). This will enable the selection of safe, stable and effective live attenuated RSV vaccine candidates.

30

## 10. GENERATION OF HUMAN RESPIRATORY SYNCYTIAL VIRUS VACCINE (RSV) CANDIDATE BY DELETING THE VIRAL SH AND M2ORF2 GENES

### 10.1. M2-2 DELETION MUTANT

35 To delete M2-2 genes, two Hind III restriction enzyme sites were introduced at RSV nucleotides 8196 and 8430, respectively, in a cDNA subclone pET(S/B) which contained an

RSV restriction fragment from 4478 to 8505. The RSV restriction fragment had been previously prepared by Quikchange site-directed mutagenesis (Stratagene, La Jolla, CA). Digestion of pET(S/B) with Hind III restriction enzyme removed a 234 nucleotide sequence which contained the majority of the M2-2 open reading frame. The nucleotides encoding the first 13 amino acids at the N-terminus of the M2-2 gene product were not removed because this sequence overlaps M2-1. The cDNA fragment which contained M2-2 gene deletion was digested with SacI and BamHI and cloned back into a full-length RSV cDNA clone, designated pRSVΔM2-2

Infectious RSV with this M2-2 deletion was generated by transfecting pRSVΔM2-2 plasmid into MVA-infected Hep-2 cells expressing N, P and L genes. Briefly, pRSVΔM2-2 was transfected, together with plasmids encoding proteins N, P and L, into Hep-2 cells which had been infected with a recombinant vaccinia virus (MVA) expressing the T7 RNA polymerase. Transfection and recovery of recombinant RSV was performed as follows. Hep-2 cells were split five hours or a day before the transfection in six-well dishes. Monolayers of Hep-2 cells at 70% - 80% confluence were infected with MVA at a multiplicity of infection (moi) of 5 and incubated at 35°C for 60 min. The cells were then washed once with OPTI-MEM (Life Technologies, Gaithersburg, M.D.). Each dish was replaced with 1 ml of OPTI-MEM and 0.2 ml transfection medium was added. The transfection medium was prepared by mixing 0.5 - 0.6 µg of RSV antigenome, 0.4 µg of N plasmid, 0.4 µg of P plasmid, and 0.2 µg of L plasmid in a final volume of 0.1 ml OPTI-MEM medium. This was combined with 0.1 ml of OPTI-MEM containing 10 µl lipofectACE (Life Technologies). After a 15 minute incubation at room temperature, the DNA/lipofectACE mixture was added to the cells. The medium was replaced one day later with MEM containing 2% FBS. Cultures were further incubated at 35°C for 3 days and the supernatants harvested. Three days post-transfection, 0.3 - 0.4 ml culture supernatants were passaged onto fresh Hep-2 cells and incubated with MEM containing 2% FBS. After incubation for

six days, the supernatant was harvested and the cells were fixed and stained by an indirect horseradish peroxidase method using goat anti-RSV antibody (Biogenesis) followed by a rabbit anti-goat antibody linked to horseradish peroxidase. The virus infected cells were then detected by addition of substrate chromogen (DAKO) according to the manufacturer's instructions. Recombinant RSV which contained M2-2 gene deletion was recovered from the transfected cells. Identification of rRSVΔM2-2 was performed by RT/PCR using primers flanking the deleted region. As shown in Fig. 12A, a cDNA fragment which is 234 nucleotides shorter than the wild-type RSV was detected in rRSVΔM2-2 infected cells. No cDNA was detected in the RT/PCR reaction which did not contain reverse transcriptase in the RT reaction. This indicated that the DNA product was derived from viral RNA and not from contamination. The properties of the M2-2 deletion RSV are currently being evaluated.

#### 10.2. SH DELETION MUTANT.

To delete the SH gene from RSV, a Sac I restriction enzyme site was introduced at the gene start signal of SH gene at position of nt 4220. A unique SacI site also exists at the C-terminus of the SH gene. Site-directed mutagenesis was performed in subclone pET(A/S), which contains an AvrII(2129) SacI (4478) restriction fragment. Digestion of pET(A/S) mutant with SacI removed a 258 nucleotide fragment of the SH gene. Digestion of the pET(A/S) subclone containing the SH deletion was digested with Avr II and Sac I and the resulting restriction fragment was then cloned into a full-length RSV cDNA clone. The full-length cDNA clone containing the SH deletion was designated pRSVΔSH.

Generation of the pRSVΔSH mutant was performed as described above (see 10.1). SH-minus RSV (rRSVΔSH) was recovered from MVA-infected cells that had been co-transfected with pRSVΔSH together with N, P and L expression plasmids. Identification of the recovered rRSVΔSH was performed by RT/PCR using a pair of primers which flanked the SH gene. As

shown in Fig. 12A, a cDNA band which is about 258 nucleotides shorter than the wild-type virus was detected in the rRSVASH infected cells. No DNA was detected in the RT/PCR reaction which did not have reverse transcriptase in the RT reaction.

- 5 This indicated that the PCR DNA was derived from viral RNA and was not artifact, and that the virus obtained was truly SH-minus RSV.

### 10.3. GENERATION OF BOTH SH AND M2-2 DELETION MUTANT.

- 10 Both SH and M2-2 genes were deleted from the full-length RSV cDNA construct by cDNA subcloning. A Sac I to Bam HI fragment containing M2-2 deletion removed from cDNA subclone pET(S/B) $\Delta$ M2-2RSV was cloned into pRSV $\Delta$ SH cDNA clone. The double gene deletion plasmid pRSV $\Delta$ SH $\Delta$ M2-2 was confirmed by  
15 restriction enzyme mapping. As shown in Fig. 12B, the SH/M2-2 double deletion mutant is shorter than the wild-type pRSV cDNA.

- Recovery of infectious RSV containing both the SH and M2-2 deletion was performed as described earlier. Infectious  
20 virus with both SH and M2-2 deleted was obtained from transfected Hep-2 cells.

Table II CAT Expression levels of Mutant RSV L-gene

25	Mut.	Conc. of CAT (ng/mL)		Charge Cluster	Charge to Alanine Change	Rescued Virus
		33°C	39°C			
30	A33	0.246	Bkg	5	135E, 136K	No
	A73	3.700	0.318	3	146D, 147E, 148 D	Yes
	A171	3.020	Bkg	3	157K, 158D	Yes
	A81	1.000	0.280	3	255H, 256K	Yes
	A185	Bkg	Bkg	3	348E, 349E	No
	A91	Bkg	Bkg	3	353R, 355R	No
35	A101	Bkg	Bkg	3	435D, 436E, 437R	No
	A192	1.960	Bkg	3	510E, 511R	Yes
	A11	0.452	Bkg	1	520R	Yes



	Mut.	Conc. of CAT (ng/mL)		Charge Cluster	Charge to Alanine Change	Rescued Virus
		33°C	39°C			
5	A111	2.320	0.267	4	568H, 569E	Yes
	A121	0.772	Bkg	2	587L, 588R	No
	A133	Bkg	Bkg	4	620E, 621R	No
	A141	2.800	Bkg	3	1025K, 1026D	Yes
10	A25	0.169	Bkg	3	1033D, 1034D	Yes
	A45	5.640	0.478	5	1187D, 1188K	Yes
	A153	4.080	0.254	5	1187D, 1188K, 1189R, 1190E	Yes
	A162	10.680	Bkg	3	1208E, 1209R	No
15	A201	Bkg	Bkg	3	1269E, 1270K	No
	A211	2.440	0.047	3	1306D, 1307E	Yes
	A221	0.321	Bkg	3	1378D, 1379E	No
	A231	Bkg	Bkg	3	1515E, 1516K	No
20	A241	1.800	0.308	3	1662H, 1663K	Yes
	A57	5.660	0.706	3	1725D, 1726K	Yes
	A65	3.560	0.168	2	1957R, 1958K	Yes
	A251	0.030	Bkg.	3	2043D, 2044K	Yes
	A261	Bkg	Bkg	3	2102K, 2103H	No
25	AD11	2.800	0.456	5 and 3	1187D, 1188K, 1725D, 1726K	No
	AD21	2.640	0.226	5 and 2	1187D, 1188K, 1957R, 1958K	No
	AD31	1.280	0.192	3 and 2	1725D, 1726K, 1957R, 1958K	No
30	F1	Bkg	Bkg	-	521 F to L	Yes
	F13	0.13	Bkg	-	521 F to L	Yes
	Lwt	3.16	-	-	no amino acid changes	Yes

35 The present invention is not to be limited in scope by the specific embodiments described which are intended as

single illustrations of individual aspects of the invention,  
and any constructs, viruses or enzymes which are functionally  
equivalent are within the scope of this invention. Indeed,  
various modifications of the invention in addition to those  
5 shown and described herein will become apparent to those  
skilled in the art from the foregoing description and  
accompanying drawings. Such modifications are intended to  
fall within the scope of the appended claims.

Various publications are cited herein, the disclosures of  
10 which are incorporated by reference in their entireties.

15

20

25

30

35

The present invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and any constructs, viruses or enzymes which are functionally  
5 equivalent are within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to  
10 fall within the scope of the appended claims.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

15

20

25

30

35

WHAT IS CLAIMED IS:

1. An isolated infectious Respiratory Syncytial Viral (RSV) particle which comprises an RSV antigenome or  
5 genome containing at least one lethal deletion in the M2-ORF1 gene.
2. An isolated infectious RSV particle which  
comprises a chimeric RSV antigenome or genome encoding  
10 antigenic polypeptides of both RVS-A and RSV-B.
3. An isolated infectious RSV particle having an  
attenuated phenotype comprising an RSV antigenome or genome  
containing an L gene mutation.  
15
4. The isolated infectious RSV particle of Claim  
1, 2, or 3 which further comprises a heterologous sequence.
5. The isolated infectious RSV particle of Claim 4  
20 in which the heterologous sequence is derived from the genome  
of influenza.
6. A recombinant RNA molecule comprising a binding  
site specific for a RSV RNA-directed RNA polymerase of a  
25 negative strand RNA virus operatively linked to a RSV RNA  
containing a deletion in M2-ORF1 or M2-ORF2 and further  
containing a heterologous RNA sequence comprising the reverse  
complement of a coding sequence.
- 30 7. The recombinant RNA molecule of Claim 6 in  
which the heterologous sequence is derived from the genome of  
a virus other than RSV.
8. The recombinant RNA molecule of Claim 6 in  
35 which the heterologous sequence is derived from the genome of  
another strain of RSV.

9. The recombinant RNA molecule of Claim 8 in which the heterologous coding sequence encodes G or F gene products.
- 5 10. The recombinant RNA molecule of Claim 6 which further comprises a mutation in the L gene.
11. The recombinant RNA molecule of Claim 6 which further comprises a mutation in the SH gene.
- 10 12. A method of producing a chimeric RSV virus, comprising culturing a host cell containing nucleotide sequences encoding the N, P and L gene products of RSV and the RSV antigenome or genome in the absence of expression of the
- 15 RSV M2-ORF.
13. A vaccine comprising a chimeric RSV the genome of which contains the reverse complement of an mRNA coding sequence operatively linked to a polymerase binding site of an
- 20 RSV and a pharmaceutically acceptable carrier.
14. The vaccine of Claim 13 in which the mRNA coding sequence encodes a mutated SH gene.
- 25 15. The vaccine of Claim 13 in which the mRNA coding sequence encodes a mutated L gene.
16. The vaccine of Claim 13 in which the mRNA coding sequence encodes a mutated NS1 gene.
- 30 17. The vaccine of Claim 13 in which the mRNA coding sequence encodes a mutated M2 gene.
18. The vaccine of Claim 13 in which the mRNA
- 35 coding sequence encodes G and F genes of both RSV A and B.

19. The vaccine of Claim 13 which encodes a heterologous gene.

20. The vaccine of Claim 19 in which the heterologous gene is derived from the genome of influenza.

21. An attenuated genetically engineered RSV containing at least one modified viral gene sequence so at least some defective particles are produced during each round of viral replication in a host.

22. The attenuated virus of Claim 21 in which the sequence modified is a non-coding region that results in down-regulation of synthesis of a viral gene.

23. The attenuated virus of Claim 21 in which the sequence modified gene sequence encodes at least one insertion, deletion, or substitution of an amino acid residue or epitope.

24. A pharmaceutical composition comprising the attenuated phenotype of Claim 22 or 23.





FIG. 2







FIG. 3


## A. RSVB-GF



## B. RSVB9320G-F/M2



## C. RSVB9320G-SH/G



FIGS. 4A-C





FIG. 5






FIG. 7




FIG. 8




FIG. 9






MDPIINGNSANVYLT	DSYIKGVTSFSCNA	LGSYIFNGPIYKNDY	TNLSIQNPFLIEHNN	LKKLNTQTSLSKXV	75
KGEIKLEETPYQSL	LMTYKSTSTSEQIAT	TNLIKTIIRRAIEIS	DVKVYAIANKLGKE	KDKIKSNNGQEDNS	150
VITTIKIDILSAVK	DNQSELKADRNHSTK	QKDTIKTTLKLMC	SMQHPSPSWLHWFNL	YTKLNLILQYRSNE	225
VKNHGFTLIDNQTL	GFQFTINQYGVCIYH	KELKRIITVTYNQFL	TWMDISLIRNVCCLI	TWISNCINTLKSIG	300
LRCGFNNVILITQFL	YGDCELLKLFHNEGFI	IIKEVEGFIMSLIN	ITWEDQFRKRPVNSM	LNNITDAANKAKQNL	375
LSRVCHTLIDKVTSL	NILNGRWTLLSKFL	KLKLAGDNLNINLS	ELYFLPFIIFGHMVD	ERQAMDVAKINQNET	450
RFYLLSSLSMGRCAF	YIRIKTGFGVNNYRW	PTLRNAIVLPURMLT	YKYNITVPSLELLETE	RDLIVLSGLRFRFREF	525
RLPKVKVLEIMINDK	AISPPKNLITWTFPR	NYMPSHIQNYTEHEK	LKFSSESKSRVLEY	YLRDNKFNEDCLYNC	600
VNQSYLNPNHNVVS	LTGKERELSVGRMFA	MQPMFRQVQILAEK	MTAENILQFFPESLT	RYGDLELQKILELKA	675
GTISNKSNNYNDVNN	YTSKRSITLDSKFN	QAFRYETSCISDVL	DELHGVOSLFSWLHL	TIPHTVITICTYRHAP	750
PYIGDHIVDLNNVDE	QSGLYRYHMGGIEGW	EQKLTWTEAISLIDL	ISLKGKESIATALING	DNQSIDISKPIRLME	825
GOTHAQADVLLAINS	LKLAJKEVAGIGHKL	KGTETYISRDQMOMS	KTQHQNVVYYPASIK	KVLRVGFWINITLDD	900
FKVLSLESTGSLQDEL	EYRGESLLCSLIFRN	VMLNQINQALQKNSA	LCNNKLYLDILKVLK	HLKTFPFLNMDITDL	975
TLXNMLPLFGGGDP	NLLYRSFYTTRTPDFL	TEALVHSFYVILSYTT	NHDLKKQLQDLSDDR	LKNFLTCIITFDKNP	1050
NABFTVLMRDPQALG	SERQAKITSEINRLA	VTEVLSTAPAKTESK	SAQHYTTTETDLNDI	MONIBETYPHGLRW	1125
YESLFPYKAEKIVNL	ISGTSKSTINILEKTS	AIDITDIDDRATEMMR	KNTILLIRIILPDQN	RKRREILSMENLSIT	1200
ELSKVVRERSWLSN	IVGVTPSPMTYMTDI	KYTTSTISSGIIIEK	YVNSLSLTRGERGPTK	PWVGSTQKCKTWPV	1275
YNRQVLTGKQROQID	LLAKLDWVYASTDNK	DEFMBEELSITGILGT	YEKAKXLFPOYLSVN	YLRULTVSVSRPQREF	1350
ASIPAVRTYNNYHPTD	SPINRILTEKYGED	IDIVFQNCISFGLSL	MSVVEQFTNVCPNRI	ILIPKLINEIHLAKPP	1425
IFPTGDVHHKLVQVI	QKQNFPLPKXISLTQ	YVEFLFNKNTKLSGS	HYNSNLIHLAKISDY	FHTNTYILSTNLAGHW	1500
ILITIQMDKSKGLIFE	KDMGEGYITDHFIFIN	LKVFNFNAKYTVLLFC	HKGYGKAKLECDMNT	SDLLCVLELIDSSVW	1575
KESMKVFLQKVTKY	ILSQDASLHRVKGCH	SEKLMELKRLANVAEF	TGCPWVNVNIDYHPTH	MKAILTYVIDLVRMGL	1650
INIDRTHIRNKHGFN	DEFTYNTYININYNF	SDNTHLITKHLIRIAN	SELENNVNDYHPTP	ETLENILANPIKSDN	1725
KKTLINDYCICGVNDS	IMPLPLSNKLIKSSS	AMIRNTYNSKQDLXAN	FPMAVVIDRIIDHSGN	TAKSQOLYTTTSHQI	1800
SLVHNSYTSICMLFW	HHINRNFNFVSSTGC	KISIEYILKDLKIKD	PNCIAFTIGEGAGNLL	LRTVVELHLPDIRIYI	1875
RSIAKQDNHDSLPTER	LRLXNGHINIDYGEN	LTIIPATDATNNIHWIS	YLHFKAPBETISLFC	DEANLSTVANNKIIII	1950
EMSKHVKKKQYCSVV	NKCMILIVYKHAQDDI	DFKLDNITILTKTYVC	LGSKLSKSEVVIIVLT	IGCAPLFPFVNNVQIN	2025
AKLLISRTKNFIMPK	KADKESIDANIKKTSI	PFLCYPTKKGINTA	LSKLSKSVSGDILSY	STAGRNVEVFSNKLIN	2100
HKHMNLIKWFNNHVLN	FRSTELATNHLVWVE	STVPYLSLANSILTT	NELKLIKIKITGSLIY	NFHNE	2165

C Cysteine residues

C Cysteine residues that were changed to valine or aspartic acid

C Cysteine residue deleted

FIG. 11



FIGS. 12A-B

# DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below at 201 et seq. underneath my name.

I believe I am the original, first and sole inventor if only one name is listed at 201 below, or an original, first and joint inventor if plural names are listed at 201 et seq. below, of the subject matter which is claimed and for which a patent is sought on the invention entitled

## RECOMBINANT RSV EXPRESSION SYSTEMS AND VACCINES

and for which a patent application:

☐ is attached hereto and includes amendment(s) filed on \_\_\_\_\_ (if applicable)  
☐ was filed in the United States on \_\_\_\_\_ as Application Serial No. \_\_\_\_\_ (for declaration not accompanying application)  
with amendment(s) filed on \_\_\_\_\_ (if applicable)  
☐ was filed as PCT international application Serial No. \_\_\_\_\_ on \_\_\_\_\_ and was amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified application, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED PRIOR TO THE FILING DATE OF THE APPLICATION			
APPLICATION NUMBER	COUNTRY	DATE OF FILING (day, month, year)	PRIORITY CLAIMED
			YES <input type="checkbox"/> NO <input type="checkbox"/>
			YES <input type="checkbox"/> NO <input type="checkbox"/>

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

APPLICATION NUMBER	FILING DATE
60/069,153	September 26, 1997
60/084,133	May 1, 1998
60/089,207	June 12, 1998

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	FILING DATE	STATUS		
		PATENTED	PENDING	ABANDONED

POWER OF ATTORNEY: As a named inventor, I hereby appoint S. Leslie Mirock (Reg. No. 18872), Harry C. Jones, III (Reg. No. 20280), Berj A. Terzian (Reg. No. 20060), Gerald J. Fintoft (Reg. No. 20822), David Weild, III (Reg. No. 21094), Jonathan A. Marshall (Reg. No. 24614), Barry D. Rein (Reg. No. 22411), Stanton T. Lawrence, III (Reg. No. 25736), Francis E. Morris (Reg. No. 24615), Charles E. Miller (Reg. No. 24576), Gidon D. Stern (Reg. No. 27469), John J. Lauter, Jr. (Reg. No. 27814), Brian M. Poissant (Reg. No. 28462), Brian D. Coggio (Reg. No. 27624), Rory J. Radding (Reg. No. 28749), Stephen J. Harbulak (Reg. No. 29166), Thomas E. Friebe (Reg. No. 29258), Laura A. Coruzzi (Reg. No. 30742), Jennifer Gordon (Reg. No. 30753), Allan A. Fanucci (Reg. No. 30256), Geraldine F. Baldwin (Reg. No. 31232), Victor N. Balancia (Reg. No. 31231), Albert P. Halluin (Reg. No. 25227), Samuel B. Abrams (Reg. No. 30605), Steven I. Wallach (Reg. No. 35402), Paul J. Zegger (Reg. No. 33821), Edmond R. Bannon (Reg. No. 32110), Bruce J. Barker (Reg. No. 33291), all of Pennie & Edmonds LLP, whose addresses are 1155 Avenue of the Americas, New York, New York 10036, 1667 K Street N.W., Washington, DC 20006 and 2730 Sand Hill Road, Menlo Park, CA 94025, and James C. Haight (Reg. No. 25,588), Gloria H. Richmond (Reg. No. 30,416), Robert Benson (Reg. No. 33,612), Jack Spiegel (Reg. No. 34,477), Susan S. Rucker (Reg. No. 35,762), David R. Sadowski (Reg. No. 32,808), Laurence J. Hyman (Reg. No. 35,551), Steven M. Ferguson (Reg. No. 38,448), Stephen L. Finley (Reg. No. 36,357), Cindy K. Fuchs (Reg. No. 38,144), John Peter Kim (Reg. No. 38,514) whose addresses are OTT/NIH, 6011 Executive Blvd., Rockville MD 20852, and each of them, my attorneys, to prosecute this application, and to transact all business in the Patent and Trademark Office connected therewith.

**SEND CORRESPONDENCE TO:** PENNIE & EDMONDS LLP  
1155 AVENUE OF THE AMERICAS  
NEW YORK, N.Y. 10036-2711

**DIRECT TELEPHONE CALLS TO:**  
PENNIE & EDMONDS LLP DOCKETING  
(212) 790-2803

2 0 1	FULL NAME OF INVENTOR	LAST NAME Jin	FIRST NAME Hong	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY Sunnyvale	STATE OR FOREIGN COUNTRY California	COUNTRY OF CITIZENSHIP People's Republic of China	
	POST OFFICE ADDRESS	STREET 354 Connemara Way	CITY Sunnyvale	STATE OR COUNTRY California	ZIP CODE 94087
2 0 2	FULL NAME OF INVENTOR	LAST NAME Tang	FIRST NAME Roderick	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY San Carlos	STATE OR FOREIGN COUNTRY California	COUNTRY OF CITIZENSHIP Malaysia	
	POST OFFICE ADDRESS	STREET 730 Chestnut Street, #3	CITY San Carlos	STATE OR COUNTRY California	ZIP CODE 94070
2 0 3	FULL NAME OF INVENTOR	LAST NAME Li	FIRST NAME Shengqiang	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY Cupertino	STATE OR FOREIGN COUNTRY California	COUNTRY OF CITIZENSHIP People's Republic of China	
	POST OFFICE ADDRESS	STREET 21001 Seven Springs Parkway	CITY Cupertino	STATE OR COUNTRY California	ZIP CODE 94043
2 0 4	FULL NAME OF INVENTOR	LAST NAME Bryant	FIRST NAME Marty	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY Los Altos	STATE OR FOREIGN COUNTRY California	COUNTRY OF CITIZENSHIP U.S.A.	
	POST OFFICE ADDRESS	STREET 1664 Clay Drive	CITY Los Altos	STATE OR COUNTRY California	ZIP CODE 94024

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201  _____ Hong Jin	SIGNATURE OF INVENTOR 202  _____ Roderick Tang	SIGNATURE OF INVENTOR 203  _____ Shengqiang Li
DATE	DATE	DATE
SIGNATURE OF INVENTOR 204  _____ Marty Bryant	SIGNATURE OF INVENTOR 205  _____	SIGNATURE OF INVENTOR 206  _____
DATE	DATE	DATE